

## Alcohol and violence: neuropeptidergic modulation of monoamine systems

Klaus A. Miczek,<sup>1,2</sup> Joseph F. DeBold,<sup>2</sup> Lara S. Hwa,<sup>2</sup> Emily L. Newman,<sup>2</sup> and Rosa M. M. de Almeida<sup>3</sup>

<sup>1</sup>Departments of Pharmacology, Psychiatry, and Neuroscience, Tufts University, Boston, Massachusetts. <sup>2</sup>Department of Psychology, Tufts University, Medford, Massachusetts. <sup>3</sup>Department of Psychology, LPNeC, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

Address for correspondence: Klaus A. Miczek, Department of Psychology, Tufts University, 530 Boston Ave (Bacon Hall), Medford, MA 02155. klaus.miczek@tufts.edu

Neurobiological processes underlying the epidemiologically established link between alcohol and several types of social, aggressive, and violent behavior remain poorly understood. Acute low doses of alcohol, as well as withdrawal from long-term alcohol use, may lead to escalated aggressive behavior in a subset of individuals. An urgent task will be to disentangle the host of interacting genetic and environmental risk factors in individuals who are predisposed to engage in escalated aggressive behavior. The modulation of 5-hydroxytryptamine impulse flow by gamma-aminobutyric acid (GABA) and glutamate, acting via distinct ionotropic and metabotropic receptor subtypes in the dorsal raphe nucleus during alcohol consumption, is of critical significance in the suppression and escalation of aggressive behavior. In anticipation and reaction to aggressive behavior, neuropeptides such as corticotropin-releasing factor, neuropeptide Y, opioid peptides, and vasopressin interact with monoamines, GABA, and glutamate to attenuate and amplify aggressive behavior in alcohol-consuming individuals. These neuromodulators represent novel molecular targets for intervention that await clinical validation. Intermittent episodes of brief social defeat during aggressive confrontations are sufficient to cause long-lasting neuroadaptations that can lead to the escalation of alcohol consumption.

**Keywords:** violence; aggression; alcohol; GABA; glutamate; serotonin

### Introduction

Alcohol has been linked to at least half of all violent assaults, cases of child abuse, and other incidents of domestic violence as well as homicides and murders.<sup>1–3</sup> This association has persisted over several decades and extends across most regions of the world. While the consumption of alcohol features prominently in many celebratory and salient events in social life, it is the antisocial, destructive nature of alcohol's effects that is of profound concern to the criminal justice system, and as a public health problem.<sup>4,5</sup> Surprisingly, the neurobiological processes underlying the link between alcohol and social, aggressive and violent behavior remain poorly understood, and there is no effective clinical treatment for violence induced or heightened by alcohol.

Initially, this review highlights several fundamental and essential pharmacological features of the link between alcohol and aggression, ranging from the acute effects to the effects of withdrawal after long-term use. Next, we consider the reinforcing actions of aggression and alcohol, both acting on neural circuits in mesolimbic monoaminergic systems. Furthermore, this review links alcohol and aggression to a postulated deficiency in brain serotonin and will outline the different serotonin subsystems and their receptor subtypes. Serotonin (5-hydroxytryptamine, 5-HT) and other canonical monoaminergic pathways are modulated by gamma-aminobutyric acid (GABA) and glutamate, and these excitatory and inhibitory amino acids are critical sites of action for alcohol to escalate aggressive behavior. Finally, we consider some

of the prominent neuropeptides and neuroactive steroids and their modulatory action on alcohol and aggression.

There are several new conclusions about the neurobiological processes that mediate the effects of alcohol on aggressive behavior:

- (1) Significant aggression-heightening effects are evident after acute, low alcohol doses as well as during withdrawal from long-term alcohol use; alcohol may lead to the distortion of social signals from benign to threatening.
- (2) A significant subset of individuals engages in alcohol-heightened aggression, while most alcohol-consuming individuals do not exhibit significant changes in aggressive behavior.
- (3) Subsystems of 5-HT neurons originating in the dorsal raphe nuclei and maturing during specific critical periods appear to contribute to the emergence of aggressive behavior.
- (4) The modulation of 5-HT impulse flow by GABA and glutamate, acting via distinct ionotropic and metabotropic receptor subtypes, is of critical significance in the suppression and escalation of aggressive behavior.
- (5) In anticipation and reaction to aggressive behavior, neuropeptides such as corticotropin-releasing factor (CRF), neuropeptide Y (NPY), opioid peptides, and vasopressin interact with monoamines, GABA, and glutamate to attenuate and amplify aggressive behavior. These neuromodulators represent novel targets for intervention and await clinical validation.
- (6) Intermittent episodes of brief social defeat during aggressive confrontations are sufficient to cause long-lasting neuroadaptations that lead to the escalation of alcohol consumption. Activation of the CRF system is necessary for this social stress-induced escalation of alcohol consumption.

## Epidemiological background

A consistent series of epidemiological and criminal statistics documents the association between alcohol and violent crimes starting in the 1950s and extending to the present (Refs. 6–11 and Table 1). Clinical diagnoses distinguish among instrumental, calculating, premeditated, and proactive “cold” aggressive acts on the one hand,

and hostile, affective, impulsive, and reactive “hot” aggression on the other.<sup>12–14</sup> Alcohol profoundly exacerbates the latter subtypes of aggression. Important features of the epidemiological data link alcohol and several major types of violent behavior ranging from homicides, sexual assaults, rapes, and child abuse to other physical assaults. This link is large in magnitude and costly to the individual, family, and society.<sup>15–18</sup> In addition to the pervasive escalation of many types of impulsive and reactive violent acts, possibly due to the myopic perception of threatening stimuli,<sup>19</sup> some epidemiological data point to geographic hot spots and epidemics of the alcohol-violence link.<sup>20</sup> Rather than proposing a neurogenetic deterministic account of the genesis of alcohol-heightened aggressive behavior, this evidence points to several risk factors that may interact with the social context.<sup>21,22</sup>

## Discovery of ethanol’s action on discrete receptors in the brain: 1980–1990

A major shift in research on neural sites and mechanisms of ethanol and by implication on the aggression effects of ethanol began in the 1980s when the focus on the disordering effects of alcohol was redirected from the lipid component of cell membranes to the action of ethanol on protein components in neuronal membranes.<sup>23</sup> This latter action was identified at several ionotropic receptors at millimolar concentrations that were more relevant to the excitatory effects of ethanol (Fig. 1).

GABA<sub>A</sub> receptors were among the first ionotropic receptors shown to increase chloride ion (Cl<sup>-</sup>)-flux as a result of action by 20–100 mM ethanol, and this effect was blocked by GABA<sub>A</sub> antagonists.<sup>24–27</sup> GABA<sub>A</sub> receptors are pentameric ligand-gated ion channel complexes composed of at least 16 subunits, with two alpha and beta subunits plus either a gamma or delta subunit being the most typical compositions for receptors in the mammalian central nervous system (CNS).<sup>28</sup> With the aid of molecular genetic techniques, it appears feasible to learn which subunits need to be expressed in specific brain regions for alcohol to achieve specific behavioral effects. For example, GABA<sub>A</sub> receptors that contain a delta subunit instead of a gamma subunit are sensitive to concentrations of ethanol that are in the mild to moderately intoxicating range<sup>29,30</sup> (but see Ref. 31).

**Table 1.** Estimates of association between alcohol consumption and homicide rates, hazardous drinking pattern score, and estimated fraction of homicides attributable to alcohol consumption<sup>3</sup>


Author(s)	Country/area, period	Parameter estimate (SE)	Level of hazardous drinking pattern	Attributable fraction (AF)
Rossow <sup>275</sup>	North Europe, 1950–1995	0.124 (0.038)	High	0.50
Rossow <sup>275</sup>	Central Europe, 1950–1995	0.085 (0.023)	Medium	0.55
Rossow <sup>275</sup>	South Europe, 1950–1995	0.055 (0.017)	Low	0.61
Rossow <sup>276</sup>	Canada, Ontario, 1950–1995	0.093 (0.040)	Medium	0.58
Rossow <sup>276</sup>	Canada, Quebec, 1950–1995	−0.030 (0.077) ns	Low	na
Bye <sup>277</sup>	Belarus and Russia, 1959–2004	0.072 (0.016)	Very high	0.57
Bye <sup>277</sup>	Former Czechoslovakia, 1953–1989	0.117 (0.067)	Medium	0.73
Landberg and Norström <sup>278</sup>	United States, 1950–2002	0.094 (0.044)	Medium	0.57
Landberg and Norström <sup>278</sup>	Russia, 1959–1998	0.081 (0.015)	Very high	0.73
Norström <sup>279</sup>	United States, 1950–2002, dry	0.035 (0.047) ns	Low	na
Norström <sup>279</sup>	United States, 1950–2002, moderate	0.071 (0.037)	Medium	0.51
Norström <sup>279</sup>	United States, 1950–2002, wet	0.174 (0.045)	High	0.80
Ramstedt <sup>11</sup>	Australia, 1950–2003	0.075 (0.028)	Medium	0.56

ns, not significant; na, not applicable.

While ethanol has enhancing effects on GABAergic synapses, it has the opposite effect on glutamatergic signaling. Ethanol at concentrations of 10–25 mM inhibits *N*-methyl-D-aspartic acid (NMDA) glutamate receptors.<sup>32–35</sup> Intoxicating doses of ethanol can reduce glutamatergic neurotransmission in the nucleus accumbens (NAcc)<sup>36</sup> and amygdala.<sup>37</sup>

NMDA receptors (NMDARs) are tetrameric ligand-gated ion channel complexes composed of two GluN1 subunits and two GluN2 (A, B, C, or D) subunits; GluN3 (A and B) subunits reduce ligand sensitivity and Ca<sup>2+</sup> permeability.<sup>38,39</sup> As with the GABA<sub>A</sub> receptor, many NMDAR subunit

combinations are responsive to ethanol. The recombinant approach has demonstrated that GluN1 is required, but then it is the fundamental obligatory NMDAR subunit.<sup>40</sup> Of the four isoforms of GluN2 subunit, ethanol can inhibit recombinant GluN1/GluN2A and GluN1/GluN2B NMDARs, and those containing GluN2B have been found to be the most sensitive to ethanol in HEK cells, but not *Xenopus* oocyte expression systems.<sup>41–45</sup> Ethanol sensitivity is lower if the recombinant receptors contain GluN2C or GluN2D subunits instead of GluN2A or GluN2B.<sup>46</sup> The conditional knockout approach suggests that GluN2B subunits play an important role in ethanol's effects on NMDARs



**Figure 1.** Theories of ethanol action at the cellular level. Adapted from Dunwiddie,<sup>267</sup> works cited (Refs. 26, 27, 32, 34, and 268–274).

in the bed nucleus of the stria terminalis.<sup>47</sup> In addition, acute low concentrations of ethanol alter NMDA subunit trafficking and can result in a selective internalization of GluN2A subunits, as well as a change in the GluN2A/GluN2B ratio.<sup>48</sup>

Ethanol can also increase neural activity in the ventral tegmental area (VTA) and increase dopamine (DA) release in the NAcc, potentially indicative of the rewarding effects of this drug<sup>49–52</sup> (see next). These effects appear to be mediated by ethanol's actions on GABAergic and glutamatergic neurons, and in part by ethanol's effects on other ion channels, including G protein-gated inwardly rectifying K<sup>+</sup> channels (GIRK)<sup>53,54</sup> and other intracellular signaling pathways.<sup>55</sup>


Similarly, ethanol increases 5-HT release in the NAcc,<sup>56–58</sup> and this effect may play a role in modulating DA release in this nucleus.<sup>59</sup> Some of the effects of ethanol on serotonergic neurotransmission, like those on DA, appear to be indirect.<sup>60</sup> However, ethanol does directly potentiate 5-HT<sub>3</sub> receptors,<sup>61,62</sup> and these receptors, as well some other 5-HT subtypes, may be important in alcohol abuse.<sup>63–67</sup> Genetic differences in serotonergic systems across individuals may also explain some of the differences in alcohol intake and abuse<sup>68</sup> (see next).

Recently, the action of alcohol in the dorsal raphe nucleus (DRN) has been explored using electrophysiological techniques. DRN inhibitory transmission is particularly sensitive to ethanol, and this is mediated primarily by synaptic GABA<sub>A</sub> receptors and secondarily by glycine receptors.<sup>69</sup> In mice previously exposed to chronic intermittent ethanol vapor, there was increased DRN excitability

during withdrawal<sup>70</sup> and decreased spontaneous inhibitory transmission in DRN-containing brain slices from these animals. Further, bath application of ethanol enhanced the frequency of miniature inhibitory postsynaptic currents in DRN neurons in slices from ethanol-withdrawn mice, but not ethanol-naïve mice.<sup>70</sup> Chronic, voluntary ethanol intake through intermittent access to two-bottle choice in Sprague–Dawley rats and C57BL/6J mice also can lead to functional DRN changes.<sup>71,72</sup> Repeated binge drinking during adolescence specifically decreased expression of 5-HT-related genes and of genes encoding the GABA<sub>A</sub> receptor  $\alpha$ 2,  $\alpha$ 3, and  $\alpha$ 5 subunits in the DRN.<sup>73</sup> However, this contrasts with a recent clinical study demonstrating that adult alcoholics have greater tryptophan hydroxylase 2 mRNA and protein in the DRN, compared to nonpsychiatric controls.<sup>74</sup> Taken together, these results suggest that ethanol can cause alterations in DRN signaling, both after acute and chronic exposure.

### Basic pharmacology of alcohol and aggression

It has been challenging to determine the exact amount of alcohol circulating in the blood, and the amount acting on pivotal target sites in the brain, of human perpetrators or victims of violence. Instead, estimates rely on retrospective analysis of events that occurred at varying times in the past and involve questionable recall. It is even rarer to learn about precise blood levels of alcohol at the time of apprehension of an alcohol-intoxicated perpetrator




**Figure 2.** Number of aggressive monetary subtraction responses per session after the administration of placebo or three different doses of alcohol. Data points represent means of four subjects each in the low- and high-provocation conditions, and vertical lines are  $\pm 1$  SEM. From Cherek *et al.*<sup>79</sup>

or victim of aggressive acts or both. In the absence of such data, experimental studies with human subjects, and preclinical studies in animal models, supply important information on the basic pharmacology of alcohol on aggressive behavior, although experimental work on intensely violent acts is ethically unacceptable. We have developed preclinical methods to study escalated aggression after oral consumption of alcohol in a significant subgroup of individual mice, rats, and monkeys.<sup>16,75–77</sup>

Acute exposure to alcohol results in a biphasic dose- and time-dependent relationship between alcohol and most of its behavioral, physiolog-

ical, and neurochemical effects, among them aggressive behavior.<sup>78</sup> The level of circulating alcohol is systematically related to the rate and intensity of aggressive behavior in several experimental models. Low doses of alcohol on the ascending limb of the dose–effect curve result in increased aggressive behavior, whereas high doses decrease this behavior as part of its sedative effects in humans, nonhuman primates, and rodents (Fig. 2).<sup>79–83</sup> For example, resident mice or rats that were administered ethanol by gavage showed peak escalation of attack bites and sideways threats at the 1.0 g/kg dose. Similarly, when outbred Swiss mice were



**Figure 3.** Target areas of bites inflicted by resident mice after oral self-administration of water (left) or 1 g/kg ethanol (right). From Newman *et al.* (in preparation).


conditioned to self-administer fixed doses of ethanol before confronting an intruder opponent, they engaged in peak levels of aggressive behavior at the 1.0 g/kg dose.<sup>16,84</sup> In a methodological refinement, outbred Swiss mice self-administered droplets of 6% w/v ethanol voluntarily. After accumulating 1.0 g/kg ethanol within 5 min, they confronted an intruder opponent and engaged in significantly escalated aggressive behavior. Detailed quantitative analysis of alcohol-heightened aggressive behavior revealed that resident rats fail to terminate aggressive bouts.<sup>80</sup>

Aside from increasing the rate of aggressive behavior, self-administered ethanol also intensifies aggressive acts, on the basis of detailed analysis of the wounds. Among the key criteria for modeling pathological aggression in experimental animals are the lack of threats preceding injurious attacks and the indiscriminate targets for inflicting bodily harm.<sup>85,86</sup> Slow-motion video analysis reveals that alcohol self-administering resident mice or rats aim their attack bites at vulnerable parts of the intruder’s body such as the ventrum and face<sup>87</sup> (Newman *et al.*, in preparation; Fig. 3).


Epidemiological and clinical studies primarily focus on recurrent violent incidents in individuals experiencing repeated alcohol intoxication.<sup>88</sup> It is well established that half of alcohol-dependent males who cycle through bouts of intoxication and withdrawal commit violent acts.<sup>89–92</sup> Recently, an

experimental protocol was developed to identify peak aggressive behavior in laboratory mice concurrent with glutamatergic convulsive phenomena in withdrawal from 2 months of intermittent access to 20% ethanol (w/v).<sup>77</sup> With increasing length of intermittent access to alcohol, a larger proportion of mice began to engage in aggressive behavior when withdrawing from alcohol (Fig. 4).

While the evidence from preclinical studies is valuable, being based on systematic and controlled



**Figure 4.** Proportion of mice showing aggression during 8-h withdrawal from intermittent alcohol access after 1, 4, or 8 weeks of exposure. The dashed line represents the average of water drinkers across age-matched groups. \**P* < 0.05 versus week 1; #*P* < 0.05 versus H<sub>2</sub>O. From Hwa *et al.*<sup>77</sup>



**Figure 5.** Histogram representing the proportion and individual magnitude of alcohol-heightened aggression (AHA) after gavage (A) or operant self-administration (B) of 1.0 g/kg alcohol. Dark vertical bars represent outbred CFW mice whose average frequency of attack bites after 1 g/kg alcohol exceeds their baseline levels of aggression by  $>2$  SD (AHA). Gray vertical bars represent mice whose aggressive behavior is not significantly altered by 1 g/kg alcohol (alcohol nonheightened aggression, ANA), and white vertical bars represent mice whose aggressive behavior after 1 g/kg alcohol is reduced by  $>2$  SD (alcohol-suppressed aggression, or ASA). Dotted horizontal lines represent a 95% confidence interval,  $\pm 2$  SD from average baseline. From Miczek *et al.*<sup>86</sup>

drug manipulations and measurements, it is limited by important differences in patterns of voluntary alcohol consumption, alcohol uptake, distribution and metabolism, and neurochemical targets. Animal species differ considerably with regard to the fate of alcohol in the body and the brain, and to the morphological and functional characteristics of the neural sites of action for alcohol.<sup>23</sup> When translating preclinical data to the human condition, it is essential to scale doses and time of peak effect across species very carefully.<sup>23</sup>

### Individual differences in ethanol and aggression

Individuals differ markedly as to whether the consumption of a given dose of alcohol will escalate aggressive and violent behavior.<sup>93</sup> Reliable and predictive genetic, physiological, or behavioral markers remain elusive, making it difficult to identify individuals with a predisposition for heightened aggressive and violent behavior in the early phase of ethanol action or during withdrawal. Retrospective clinical studies have associated poly-

morphisms in individuals with a propensity to engage in escalated aggressive and violent behavior. For example, genotyping the *BDNF* gene polymorphism, G196A, revealed that Japanese alcoholics with violent tendencies were more likely to be homozygous for the A allele compared to nonviolent individuals.<sup>94</sup> In a Swedish sample of adolescents with alcohol-related problem behaviors, a short variant of gene encoding MAO-A was detected, particularly when the youngsters grew up in abusive families.<sup>95</sup> Longitudinal studies with a focus on childhood abuse and continuing through adolescence into early adulthood would be particularly informative in identifying interactions between developmental and genetic risk factors that engender alcohol-related violence.

Preclinical studies with mice, rats, and monkeys identify subgroups of individuals for whom a moderate dose of alcohol reliably escalates aggressive behavior, whereas most others do not show this effect<sup>16</sup> (Fig. 5). This variability in laboratory rats and mice has led to our distinction between alcohol-heightened aggressors (AHAs)

and alcohol-nonheightened aggressors on account of the persistent, large increase in injurious attack behaviors after every exposure to a moderate 1 g/kg dose of ethanol.<sup>76,80,81</sup> So far, these subtypes have been characterized in terms of pharmacological sensitivity with serotonergic, GABAergic, and glutamatergic compounds.<sup>76,84,96–102</sup> In rhesus monkeys, intense consumption of alcohol leads to long bursts of impulsive aggressive behavior primarily in those individuals with a history of social isolation early in life.<sup>103</sup> It will be most instructive to identify biomarkers that predict whether an individual will engage in alcohol-heightened aggressive behavior.


### **Victims of aggression: escalated alcohol consumption as a result of social stress**

While the aggressogenic effects of alcohol continue to be the focus of clinical and basic research with animal models,<sup>16,104</sup> a critical role of alcohol emerges in the individual who is not the perpetrator, but the target of aggressive acts. Early preclinical studies have demonstrated how administration of ethanol to mice, rats, or monkeys alters their behavior in confrontations with an aggressive opponent.<sup>105–107</sup> The higher the dose of alcohol in the intruder mouse or rat or the subordinate monkey, the more frequently these individuals provoked attacks and threats. The perception and production of social signals and pheromones contribute to the escalation of aggressive interactions. The analysis of social cues and the interpretation and integration of these cues promises to reveal how alcohol alters provocative signals, and the response to these signals. Giancola<sup>19</sup> proposes that distorted perceptions of social signals—a type of myopia—emerge as a key determinant of alcohol-escalated aggressive behavior.

When exposed to repeated episodes of aggression and displaying salient characteristics of social stress, individuals who are the target of attacks or threats can undergo neural adaptations that cause them to consume large amounts of alcohol and to exhibit signs of increased motivation for alcohol.<sup>108,109</sup> There is considerable debate about the impact of different types of environmental stressors on alcohol consumption.<sup>110–113</sup> In order to

capture the well-documented and often reiterated link between stress and alcoholic drinking in humans, several preclinical efforts have focused on social stress.<sup>114</sup> One approach focuses on continuous subordination stress that extends over several weeks and months in mice, rats, and monkeys. Low-ranking squirrel monkeys or macaques consume more alcohol than high-ranking individuals.<sup>115,116</sup> Several studies report increases in consumption of alcohol at concentrations ranging from 2% to 20% v/v in socially subordinate members of a group of mice or rats.<sup>117,118</sup> Even though these increases in intake relative to nonstressed controls are statistically significant, they are usually insufficient to produce either intoxication or dependence.

A second approach in animal models involves repeated acute episodes of social defeat stress that result in escalated alcohol drinking.<sup>119–122</sup> When CRF receptor 1 (CRF-R1) gene knockout mice bred onto an outbred CD-1 background were subjected to brief episodes of social defeat stress on 3 consecutive days, they began to consume ~3 g/kg/day of 8% alcohol (v/v), compared to the 1 g/kg/day pre-stress baseline.<sup>119</sup> This increase manifested itself only after a 3-week interval between the social stress experience and the alcohol consumption. More recently, a series of studies identified the parameters of social defeat stress that induced escalated alcohol consumption in mice without any restriction in fluid intake and engendered a preference for alcohol relative to concurrently available water. Brief episodes of social defeat stress every day or every other day were sufficient to escalate consumption of 20% unsweetened alcohol (w/v) up to 15 g/kg/day in CFW mice and up to 30 g/kg/day in C57BL/6J mice (Fig. 6).<sup>77,120</sup> Several features of these latter studies are noteworthy: the escalated alcohol intake began after several weeks without stress and continued in the absence of any further exposure to social stress, suggesting the induction of an enduring neuroadaptation. Pharmacological antagonism and genetic deletion experiments implicate the CRF system in the induction and expression of the neuroadaptations that result from exposure to brief episodes of social defeat stress and eventually lead to persistently escalated alcohol consumption (see next).




**Figure 6.** Twenty percent ethanol intake (g/kg/day) during continuous access two-bottle choice over the course of 20 days, starting 10 days after moderate ( $n = 39$ ) or mild ( $n = 19$ ) social defeat stress (control,  $n = 29$ ). Data points are 5-day averages  $\pm$  SEM beginning on the day indicated (i.e., 25 signifies days 25–29); \*\* $P < 0.001$  compared to controls. From Norman *et al.*<sup>120</sup>

### Alcohol-heightened aggression as a source of satisfaction and pleasure: mesocorticolimbic circuits

More than 50 years ago, John Paul Scott<sup>123</sup> recognized that “aggression not only produces dangers which must be avoided, but pleasures which can be enjoyed.” Among the causative factors for alcohol-heightened aggression are the rarely investigated neural mechanisms mediating the rewarding effects of aggressive behavior.<sup>124–128</sup> By contrast, the neural mechanisms for the rewarding effects of alcohol consumption are more adequately characterized. Many facets of alcohol’s reinforcing effects have been examined in a range of species and conditions.<sup>50,129–134</sup> Individuals may prefer alcohol over other commodities, seek out the opportunity to self-administer ethanol, exert themselves to obtain alcohol, or resist the negative consequences of alcohol consumption. All of these behaviors are based on neural interactions in mesencephalic–limbic–cortical loops. These circuits include sensory components, and most alcohol is consumed in flavored concoctions; alcohol solutions affect neural circuits that regulate ingestive homeostasis of intra- and extracellular fluids and calories; most importantly, the pharmacological actions of alcohol on targets in the mesocortical and mesolimbic monoaminergic pathways appear to be necessary for its reinforcing effects. While an “alcohol receptor” has not been discovered, several ionophoric receptors in GABAergic, glutamatergic, and monoaminergic

neurons are the targets of millimolar concentrations of alcohol, as previously mentioned.<sup>23</sup> Alcohol’s reinforcing effects depend on the integrity of dopaminergic neurons in the VTA that are modulated by GABAergic and glutamatergic afferents, and that project to the core and shell of the NAcc, as well as to the medial prefrontal cortex (mPFC).<sup>135</sup> These basic networks of monoamines—excitatory and inhibitory amino acids subserving the reinforcing effects of alcohol—are modulated by a large number of neuropeptides.<sup>132,136</sup>

To what extent do the neural mechanisms mediating alcohol’s reinforcing effects overlap or interact with those that are responsible for aggressive and violent acts, which in themselves function as reinforcers? So far, only a few candidate neural mechanisms have been explored in experimental models that identify aggressive acts as reinforcers. Positive allosteric modulation by benzodiazepines and allopregnanolone (ALLO) of the GABA<sub>A</sub> receptor has proven effective in escalating the rate of responding that was reinforced by the opportunity to engage in aggressive behavior.<sup>124,137</sup> As the opportunity for aggressive behavior approached, the rate of responding accelerated and plasma levels of corticosterone were elevated; and corticosterone elevations appeared to be required for responding that is motivated by the opportunity to engage in aggressive behavior and for the escalated aggression (Fig. 7).<sup>137</sup> It is likely that the allosterically modulated GABA<sub>A</sub> receptors are altering DA and 5-HT impulse flow.



**Figure 7.** The effects of fixed interval schedule of reinforcement (FI) performance and fighting on corticosterone levels. Mean plasma corticosterone levels (ng/mL) are shown from resident mice immediately or 30 min after four different events. (A) Baseline corticosterone levels at the time of day when mice are conditioned to respond on an FI10 schedule; (B) levels after a brief attack flurry; (C) levels after completing the FI, but when no fight has occurred; and (D) levels after completing the FI and engaging in a brief fight. The open bars represent corticosterone levels when the mice have not responded on the FI10, and the filled gray bars represent levels after completion of the FI10. Vertical lines indicate  $\pm 1$  SEM; \* $P < 0.05$ . From Fish *et al.*<sup>137</sup>

Direct antagonism of DA receptor subtypes reduces the rate of responding that is reinforced by the opportunity to engage in aggressive behavior; microinjections of low doses of DA D1- or D2-like receptor antagonists into the NAcc preferentially reduced the behavioral effort to gain access to an aggressive confrontation.<sup>128</sup> It appears reasonable to suggest that alcohol-heightened aggressive behavior may require intact DA activity in the mesolimbic pathway, particularly in view of heightened release of DA in the NAcc in rats that consume alcohol and subsequently engage in escalated aggressive behavior.<sup>138</sup> Recently developed optogenetic methods promise to differentiate the aggression-specific DA cells from those that mediate other motivated behavior and motor functions.

Escalated aggressive behavior with pathological features is evident in mouse and rat strains that are selectively bred for high aggression.<sup>139</sup> So far, the reinforcing nature of pathological aggression can only be inferred from the short-latency, high-frequency, and high-intensity of persistent and injurious attacks;<sup>127</sup> more definitive information

will have to be obtained from a formal analysis of the contingencies between pathologically escalated aggressive behavior and its consequences. Most evidence for neurobiological mechanisms mediating pathologically escalated aggressive behavior has focused on tonic brain 5-HT activity and on 5-HT autoreceptors.<sup>140</sup> In the prefrontal cortex (PFC), increased 5-HT has been detected when escalated aggressive behavior is initiated, whereas termination of aggressive bouts and return to tonic levels of 5-HT is associated with reduced neural activity in the serotonergic soma of raphe cells.<sup>98,141</sup> The extent to which alcohol's violence- and aggression-heightening effects depend on serotonergic activity has been studied both in clinical and preclinical models for several decades.

### The 5-HT-alcohol link: from the deficiency hypothesis to discrete subsystems

No other neurotransmitter has been more intensively investigated than 5-HT when characterizing neural mechanisms of violence and aggression,<sup>98,140,142–144</sup> including the link between

alcohol and aggression.<sup>104,145–148</sup> Historically, the most frequently reiterated hypothesis attributed a predisposition for violent behavior to a deficiency in 5-HT in alcohol-consuming individuals, supported primarily by postmortem assays of tissue samples or lumbar cerebrospinal fluid.

Shortly after the discovery of 5-HT in the brain, the calming role of this biogenic amine was highlighted,<sup>149</sup> and then extended to aggressive individuals.<sup>150</sup> Consistent evidence in many species, ranging from invertebrates, fish, and rodents to nonhuman primates and humans, documents an effective reduction in aggressive behavior after treatment with drugs that target 5-HT<sub>1A</sub> or 5-HT<sub>2A/C</sub> receptor subtypes and transporter sites.<sup>140,151–153</sup> Conversely, monkeys and humans with the short allele of the serotonin transporter polymorphism (5HTTLPR) are characterized by heightened aggressive behavior.<sup>154,155</sup> However, like clinically employed pharmacotherapeutic treatments with potent antiaggressive effects (including antipsychotics,  $\beta$ -adrenergic blockers, steroid derivatives, and benzodiazepines), nearly all of the 5-HT<sub>1A</sub> receptor agonists, 5-HT<sub>2A</sub> receptor antagonists, and selective serotonin reuptake inhibitors decrease aggressive behavior in dose ranges that have problematic side-effect profiles.<sup>156</sup> Agonists of 5-HT<sub>1B</sub> receptors appear to result in behaviorally more selective antiaggressive effects in experimental studies with mice and humans.<sup>76,102,157–159</sup>

Research on the anatomical sites of action for alcohol's effects on escalated aggression has focused on the regions within the PFC, amygdala, septum, and hypothalamus.<sup>148,156</sup> Imaging studies in patients with intermittent explosive disorder show deficient activation of the mPFC concurrent with heightened activity in the amygdaloid complex.<sup>160–162</sup> The same regions are activated by acute alcohol intake, as indicated by converging evidence from several methodologies. Expression of immediate early genes in laboratory rodents that engage in escalated aggressive behavior identify several interconnected brain regions along the center of the neuraxis, ranging from the periaqueductal gray to the ventral tegmentum, hypothalamus, preoptic area, septum, amygdala, and PFC.<sup>141,153,163–165</sup> Expression of 5-HT receptor subtypes is reduced in the PFC of those individual mice who engage consistently in

alcohol-heightened aggression relative to those who do not.<sup>99</sup>


Intracerebral microinjection studies localized the neural pathway projecting from the DRN to the striatum and PFC as pivotal in the actions of 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors. Activation of these subtypes at the 5-HT<sub>1A</sub> somatodendritic autoreceptors, as well as pre- and postsynaptic 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors, effectively reduced alcohol-heightened aggressive behavior in mice and rats.<sup>98,101,166,167</sup> Studies with muscimol microinjections suggest that escalated aggression as a result of alcohol consumption involves GABAergic modulation of 5-HT in the DRN.<sup>98</sup> A role of glutamatergic modulation of 5-HT impulse flow to the mPFC in alcohol-heightened aggression is indicated by effects of memantine and ketamine in AHA mice (Newman *et al.*, in preparation). An emerging hypothesis regarding the mechanism for alcohol-heightened aggression focuses on subpopulations of 5-HT projections to the forebrain and their modulation by specific GABA and glutamate receptors.

### Alcohol-heightened aggression and GABA

A great deal of attention has been paid to the effects of alcohol on GABA receptors for several reasons. As discussed below, alcohol can allosterically modulate GABA<sub>A</sub> receptors but a number of effects of alcohol on the brain and behavior are similar to those of more specific drugs that act relatively specifically on GABA<sub>A</sub> receptors. For example, the sedative and hypnotic effects of alcohol are similar to those of barbiturates and benzodiazepines, and both of these classes of drugs also act on GABA<sub>A</sub> receptors as positive allosteric modulators. In addition, chronic exposure to alcohol alters the expression of GABA<sub>A</sub> receptor subunits,<sup>168–171</sup> although the expression of these subunits appears to depend on the pattern and dose of alcohol, as well as whether withdrawal occurs.

#### GABA<sub>A</sub> receptors

Alcohol can increase aggressive behavior not only in people but also in a number of laboratory animal species. However, the magnitude of this effect varies across individuals.<sup>80,172</sup> We have been investigating the likely bases for individual differences in sensitivity to the proaggressive effects of alcohol in



**Figure 8.** Attack bite frequencies in a novel environment following systemic midazolam treatment of wild-type C57BL6/J and benzodiazepine-insensitive *Gabra2* (H101R) point-mutated mice. Values are means  $\pm$  SEM; \* $P < 0.05$ , \*\* $P < 0.01$  compared to vehicle; # $P < 0.05$ , ## $P < 0.01$  compared to wild type. From Newman et al.<sup>182</sup>

laboratory rodents and it is quite clear is that alcohol generally has biphasic effects on aggressive behavior (see next). Although these drugs can be proaggressive in some individuals,<sup>173,174</sup> higher doses of benzodiazepines and GABA<sub>A</sub> agonists in conjunction with neuroleptics have an inhibitory effect, and so are often used to sedate violent patients.<sup>175–177</sup>

Benzodiazepines can act as positive modulators of GABA<sub>A</sub> receptors, but not all GABA<sub>A</sub> receptors have a benzodiazepine binding site. Benzodiazepine action requires that the GABA<sub>A</sub> receptor include a  $\gamma 2$  subunit as well as  $\alpha$  and  $\beta$  and subunits.<sup>178–180</sup> However, there is debate as to which  $\alpha$  subunit of the GABA<sub>A</sub> receptor is necessary for the benzodiazepine to exert sedative, anxiolytic, amnesic, or anticonvulsive effects or to lead to abuse.<sup>181</sup> We have examined whether GABA<sub>A</sub> receptor subunit composition might also influence the effect of alcohol on aggressive behavior. We have found that benzodiazepine antagonists, which preferentially act on GABA<sub>A</sub> receptors with  $\alpha 1$  subunits, can reduce the proaggressive effects of alcohol in mice.<sup>84</sup> In addition, point mutations to the genes encoding  $\alpha 1$  (H101R) or  $\alpha 2$  (H101R), but not  $\alpha 3$  (H126R) subunits in mice can block the aggression-heightening effects of midazolam (Fig. 8).<sup>182</sup> These findings may be useful in the eventual development of

therapies to specifically reduce the association between alcohol and violence.

Direct agonists of GABA<sub>A</sub> receptors, such as muscimol, can also influence aggressive behavior. Much of the early research on the effects of muscimol on aggressive behavior in laboratory rodents reported inhibitory effects.<sup>183–186</sup> However, when microinjected into the DRN of male mice, 0.006 nmol of muscimol potentiated aggressive behavior.<sup>98,187</sup> Importantly, this effect only occurred in mice that showed alcohol-heightened aggression.<sup>98</sup> In addition, muscimol microinjections were behaviorally ineffective if they missed the DRN. This suggests that the DRN may be a particularly important site in the link between GABA<sub>A</sub> receptors and the effects of alcohol on aggression.


Another tool that has been employed to study the possible link between alcohol-heightened aggression and GABA<sub>A</sub> receptors is the neuroactive steroid ALLO, produced both in the brain and peripherally and the most potent positive modulator of GABA<sub>A</sub> receptors known.<sup>188–193</sup> Disrupted ALLO is associated with depression, stress, and anxiety.<sup>194</sup> Alcohol can increase ALLO levels in laboratory rodents and altered brain concentrations of ALLO might be a mechanism through which alcohol exerts behavioral effects such as sedation, anxiolysis, and analgesia. Likewise, some pharmacological effects of alcohol may be caused by alcohol-induced increases in ALLO synthesis.<sup>195,196</sup> Forced-swim stress elevates ALLO in the brain,<sup>197</sup> and decrements in its biosynthesis, specifically in the basolateral amygdala, can affect aggressive behavior.<sup>198,199</sup> We have seen that moderate doses of ALLO increase aggressive behavior and this effect occurs primarily in the subset of mice that previously have been classified as AHAs.<sup>200</sup> When tested in combination with alcohol, ALLO shifted the alcohol dose–response curve to the left. In addition, higher doses of ALLO, like higher doses of alcohol, inhibit aggressive behavior. That finding is similar to ALLO reducing the frequency of attacks in isolation-enhanced aggression.<sup>201</sup> This neurosteroid appears to be stress responsive and plays a major role in regulating the hypothalamic–pituitary–adrenal axis.<sup>202</sup> It is possible that individual differences in ALLO may influence stress resilience. It will be important to tease apart the specific neural mechanisms for how alcohol and ALLO affect escalated aggressive behavior.

### GABA<sub>B</sub> receptors

In addition to the ionotropic GABA<sub>A</sub> receptor, GABA actions are also mediated by the metabotropic G protein-coupled GABA<sub>B</sub> receptor. The GABA<sub>B</sub> receptor agonist baclofen has received some attention as a possible treatment for alcohol dependence,<sup>132,203,204</sup> although some studies have not found it to be significantly better than placebo.<sup>205</sup> Baclofen also has been reported to reduce the severity of alcohol withdrawal symptoms in patients.<sup>203,206,207</sup> In rats, baclofen usually reduces aggressive behavior.<sup>185,208,209</sup> However, our research demonstrates that systemically administered baclofen can actually have a biphasic dose effect on aggression in male mice.<sup>210</sup> Microinjection of baclofen into the DRN can specifically increase aggressive behavior.<sup>98,187</sup> However, that effect does not depend on alcohol and occurs in mice independent of whether they are AHAs.

### Alcohol-heightened aggression and glutamate NMDAR

The rationale for considering glutamate and its receptors as relevant to the mechanisms mediating alcohol-heightened aggressive behavior derives from the discovery that alcohol (5–50 mM) antagonizes NMDARs, possibly via the GluN1-glycine site.<sup>34,211–214</sup> A postmortem analysis of hippocampal tissue from alcoholics and cocaine addicts revealed upregulation of *GRIN2B* (gene for GluN2B) in all individuals and specific upregulation of *GRIN2D* (gene for GluN2D) in alcoholics.<sup>215,216</sup> In addition, *GRIN2A* polymorphisms may be associated with disruptions in aversion learning and may predict the development of an alcohol use disorder.<sup>217,218</sup> Polymorphisms of *GRIN2B* may serve as an indicator for the effectiveness of acamprosate in the treatment of alcohol use disorders.<sup>219</sup> Preclinical investigations point to the importance of NMDARs in species-typical social behavior. Heterozygous deletion of the GluN1 subunit renders the receptor insensitive to the crucial coagonist, glycine and thereby reduces the proportion of functional NMDARs.<sup>220</sup> GluN1 hypomorphic mice consume more alcohol and spend less time in social interactions relative to their genetically unaltered counterparts.<sup>220,221</sup> Pharmacological blockade of NMDAR can similarly dysregulate social and agonistic behavior. When mice consume a modest dose of alcohol (1 g/kg) and are treated with the uncompetitive NMDAR



**Figure 9.** Attack bites during 6- to 8-h withdrawal from 8 weeks of intermittent alcohol access after treatment with the NMDA receptor antagonist memantine (0–30 mg/kg, i.p.). Black dots represent alcohol drinkers and white dots represent water drinkers. Values are means  $\pm$  SEM; \* $P < 0.05$  versus vehicle; # $P < 0.05$  versus H<sub>2</sub>O. From Hwa et al.<sup>77</sup>

antagonist memantine, their aggressive behavior toward a submissive male conspecific escalates.<sup>96</sup> Furthermore, when alcohol-dependent mice are challenged with a moderate dose of memantine (5 mg/kg) during withdrawal from access to alcohol following 8 weeks of access, their aggressive behavior escalates (Fig. 9).<sup>77</sup> Collectively, these findings point to the genetically or pharmacologically suppressed NMDAR as pivotal in behavioral dysregulation as indicated by increased alcohol self-administration, more intense social anxiety-like behavior, and escalated aggressive behavior.

Conversely, several findings point to the therapeutic value of NMDAR antagonists as suggested by the anticraving effects of acamprosate, memantine, and neramexane (MRZ 2/579) in rats.<sup>222,223</sup> Additionally, treatment with the uncompetitive antagonist MK-801 (dizocilpine) can increase social interactions between adolescent rats, suggesting reduced social anxiety.<sup>224</sup> Importantly, in patients diagnosed with alcohol use disorders, memantine and ketamine treatments reduce both cravings for alcohol and alcohol consumption.<sup>225–227</sup>

These divergent interactions between alcohol and NMDAR antagonists may depend on the expression of the GluN2 subunit. Receptors containing GluN2A or GluN2B are more sensitive to the inhibitory action of alcohol and exhibit regionally specific expression patterns.<sup>228</sup> GluN2A-containing NMDARs are expressed throughout the CNS of rats, while GluN2B subunits are localized preferentially

in the forebrain, specifically in the prelimbic and orbital regions of the PFC.<sup>229</sup> Glutamatergic pyramidal cells within these prefrontal cortical areas are activated during aggressive encounters in rats.<sup>230</sup> Excitatory inputs, particularly from the PFC to the DRN may facilitate the initiation and termination of aggressive behavior.<sup>231</sup> We hypothesize that dysregulation of cortical inhibition of GluN1/GluN2B-containing NMDARs may result in alcohol-induced social disinhibition and escalated aggression. Individual differences in GluN2 subunit expression may determine the behavioral effects of drugs such as ethanol and memantine in interaction with an individual's predisposition for aggressive behavior and other psychopathologies.<sup>232</sup>

### **Alcohol-heightened aggression and neuropeptides: CRF and opioid peptides**

Stress-related neuropeptide systems have been implicated in alcohol-related behaviors, most prominently CRF and urocortin, NPY, nociception/orphanin, and neurokinin.<sup>233</sup> These and other neuropeptides have been implicated in the regulation and dysregulation of social behaviors, especially vasopressin, oxytocin, and substance P.<sup>234–236</sup> Most of the evidence points to a significant role of CRF and opioid peptides in the modulation of alcohol-related aggression.

#### *Corticotropin-releasing factor*

CRF is the initial link in the cascade of neural, endocrine, and behavioral responses to stressful stimuli,<sup>237,238</sup> especially those related to prosocial and conflict situations.<sup>239,240</sup> CRF biphasically modulates aggression in various species,<sup>241,242</sup> with low doses increasing and high doses decreasing agonistic behaviors in rats.<sup>243</sup> Further, most CRF-R1 antagonists effectively reduce aggression in socially organized species.<sup>244–246</sup> One-way CRF-R1 antagonism prevents alcohol-heightened aggression is, in part, its facilitation of 5-HT impulse flow from the DRN to the mPFC.<sup>247</sup> The mPFC has emerged as a critical site through which alcohol-induced impairments can contribute to emotional outbursts and violence.<sup>126,248</sup> Alcoholic, impulsive offenders,<sup>249</sup> and perpetrators of domestic violence<sup>250</sup> are characterized by dysregulated CRF levels compared to healthy controls. Enhanced aggression, compared to species-typical aggression, may recruit stress-related circuits resulting in a loss of inhibitory control.

#### *Opioid peptides*

Brain opioid peptides acting on mu, delta, or kappa opioid receptors (MOR, DOR, KOR) are critically involved in the regulation of social behaviors and mood disorders.<sup>251,252</sup> Increased aggression after acute opiates has been documented in several species such as fish,<sup>253–255</sup> cats,<sup>256</sup> pigs,<sup>257</sup> crickets,<sup>258</sup> and mice,<sup>259</sup> which implicate endogenous opioids in social conflict. By contrast, opioid antagonists reduce alcohol-heightened aggression primarily in some individuals, possibly with a distinctive pattern of expression for genes encoding MOR, DOR, and KOR.<sup>260–262</sup> Research is limited for alcohol-heightened aggression, but alcohol interferes with endogenous opioid mechanisms, including the acute release of  $\beta$ -endorphin.<sup>263</sup> We hypothesize that endorphins acting on MOR are of particular relevance to alcohol-heightened aggression, and represent important targets for intervention. Social conflict activates antinociception through opioid mechanisms.<sup>261,264,265</sup> Opioid receptor activation in aggressive encounters may lead to exaggerated responses to threatening environmental stimuli. Naltrexone can reduce self-injurious behavior,<sup>266</sup> so the prominent interaction between alcohol and endogenous opioids may enable opioid antagonists to suppress alcohol-related aggression.

#### **Future agenda**

One of the urgent tasks will be to characterize the individuals who are predisposed to engage in alcohol-heightened aggressive behavior. To disentangle the host of interacting genetic and environmental risk factors will require a concerted effort at every level of analysis.

In preclinical studies, novel neurobiological tools such as optogenetic stimulation and inhibition of chemically specific cell groups in discrete nodes of neural microcircuits promise to characterize effective targets of intervention to attenuate alcohol-heightened aggressive behavior.

Novel molecular targets in the neural microcircuits for escalated aggressive behavior are likely to emerge from studies with neuropeptidergic modulation of GABA and glutamate cells that stimulate or inhibit monoaminergic pathways. Alternatively, it will be useful to explore steroid systems and endocannabinoids for their role in alcohol-heightened aggression.

The validity of experimental models of alcohol-heightened aggression derives from their potentially injurious and harmful nature. The ethical dilemma on the one hand of studying alcohol-escalated aggressive behavior in experimental models and on the other hand of following the principle of harm reduction will require careful and thorough examination in each case.

### Conflicts of interest

The authors declare no conflicts of interest.

### References

- Pernanen, K. 1976. "Alcohol and crimes of violence." In *Social Aspects of Alcoholism*. B. Kissin, Ed.: 351–444. New York: Plenum Press.
- Roizen, J. 1993. "Issues in epidemiology of alcohol and violence." In *Alcohol and Interpersonal Violence: Fostering Multidisciplinary Perspectives. Research Monograph No. 24*. S.E. Martin, Ed.: 3–36. Rockville, MD: National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism.
- Rossow, I. & E.K. Bye. 2013. "The problem of alcohol-related violence: an epidemiological and public health perspective." In *Alcohol-Related Violence: Prevention and Treatment*. M. McMurrin, Ed.: 1–18. Chichester: Wiley-Blackwell.
- Holder, H.D. 2008. Alcohol and violence: a complex nexus of drinking environment and drinking pattern. *Addiction* **103**: 78–79.
- Shield, K.D., C. Parry & J. Rehm. 2013. Chronic diseases and conditions related to alcohol use. *Alcohol Res.* **35**: 155–173.
- Tinklenberg, J.R. & F.M. Ochberg. 1981. "Patterns of adolescent violence: a California sample." In *Biobehavioral Aspects of Aggression*. D.A. Hamburg, Ed.: 121–140. New York: Alan R. Liss.
- Shupe, L.M. 1954. Alcohol and crime. A study of the urine alcohol concentration found in 882 persons arrested during or immediately after the commission of a felony. *J. Crim. Law Criminol Police Sci.* **44**: 661–664.
- Livingston, L.R. 1986. Measuring domestic violence in an alcoholic population. *J. Sociol. Soc. Welfare* **13**: 934–953.
- Rossow, I. 2004. Alcohol and homicide in Russia. *Addiction* **99**: 937.
- Norström, T. 2011. The role of alcohol in the Russian mortality crisis. *Addiction* **106**: 1957–1965.
- Ramstedt, M. 2011. Population drinking and homicide in Australia: a time series analysis of the period 1950–2003. *Drug Alcohol Rev.* **30**: 466–472.
- Vitiello, B. & D.M. Stoff. 1997. Subtypes of aggression and their relevance to child psychiatry. *J. Am. Acad. Child Adolesc. Psychiatry* **36**: 307–315.
- Lesch, K.P. & U. Merschdorf. 2000. Impulsivity, aggression, and serotonin: a molecular psychobiological perspective. *Behav. Sci. Law* **18**: 581–604.
- Kessler, R.C., E.F. Coccaro, M. Fava, *et al.* 2006. The prevalence and correlates of DSM-IV intermittent explosive disorder in the National Comorbidity Survey Replication. *Arch. Gen. Psychiatry* **63**: 669–678.
- Collins, J.J. & P.M. Messerschmidt. 1993. Epidemiology of alcohol-related violence. *Alcohol Health Res. World* **17**: 93–100.
- Miczek, K.A., E.W. Fish, R.M.M. de Almeida, *et al.* 2004. Role of alcohol consumption in escalation to violence. *Ann. N. Y. Acad. Sci.* **1036**: 278–289.
- Smith, G.S., C.C. Branas & T.R. Miller. 1999. Fatal nontraumatic injuries involving alcohol: a metaanalysis. *Ann. Emerg. Med.* **33**: 659–668.
- Egami, Y., D.E. Ford, S.F. Greenfield & R.M. Crum. 1996. Psychiatric profile and sociodemographic characteristics of adults who report physically abusing or neglecting children. *Am. J. Psychiatry* **153**: 921–928.
- Giancola, P.R. 2013. "Alcohol and aggression: theories and mechanisms." In *Alcohol-Related Violence: Prevention and Treatment*. M. McMurrin, Ed.: 37–59. Chichester: John Wiley & Sons.
- Cortez, F.C.P. & D.A. Baltieri. 2013. "Alcohol-related violence: an international perspective." In *Alcohol-Related Violence: Prevention and Treatment*. M. McMurrin, Ed.: 19–33. Chichester: John Wiley & Sons.
- Gruenewald, P.J., L. Remer & R. Lipton. 2002. Evaluating the alcohol environment: community geography and alcohol problems. *Alcohol Res. Health* **26**: 42–48.
- Gorman, D.M., P.J. Gruenewald & L.A. Waller. 2013. Linking places to problems: geospatial theories of neighborhoods, alcohol and crime. *GeoJournal* **78**: 417–428.
- Tabakoff, B. & P.L. Hoffman. 2013. The neurobiology of alcohol consumption and alcoholism: an integrative history. *Pharmacol. Biochem. Behav.* **113**: 20–37.
- Mehta, A.K. & M.K. Ticku. 1988. Ethanol potentiation of GABAergic transmission in cultured spinal cord neurons involves gamma-aminobutyric acidA-gated chloride channels. *J. Pharmacol. Exp. Ther.* **246**: 558–564.
- Allan, A.M. & R.A. Harris. 1986.  $\gamma$ -Aminobutyric acid and alcohol actions: neurochemical studies of long sleep and short sleep mice. *Life Sci.* **39**: 2005–2015.
- Suzdak, P.D., R.D. Schwartz, P. Skolnick & S.M. Paul. 1986. Ethanol stimulates  $\gamma$ -aminobutyric acid receptor-mediated chloride transport in rat brain synaptosomes. *Proc. Natl. Acad. Sci. U.S.A.* **83**: 4071–4075.
- Ticku, M.K., P. Lowrimore & P. Lehoullier. 1986. Ethanol enhances GABA-induced  $^{36}\text{Cl}^{-}$ -influx in primary spinal cord cultured neurons. *Brain Res. Bull.* **17**: 123–126.
- Baur, R., F. Minier & E. Sigel. 2006. A GABA<sub>A</sub> receptor of defined subunit composition and positioning: concatenation of five subunits. *FEBS Lett.* **580**: 1616–1620.
- Hanchar, H.J., M. Wallner & R.W. Olsen. 2004. Alcohol effects on  $\gamma$ -aminobutyric acid type A receptors: are extrasynaptic receptors the answer? *Life Sci.* **76**: 1–8.
- Wallner, M., H.J. Hanchar & R.W. Olsen. 2003. Ethanol enhances  $\alpha_4\beta_3\delta$  and  $\alpha_6\beta_3\delta$   $\gamma$ -aminobutyric acid type A receptors at low concentrations known to affect humans. *Proc. Natl. Acad. Sci. U.S.A.* **100**: 15218–15223.
- Borghese, C.M., S. Storustovu, B. Ebert, *et al.* 2006. The  $\delta$  subunit of  $\gamma$ -aminobutyric acid type A receptors does not confer sensitivity to low concentrations of ethanol. *J. Pharmacol. Exp. Ther.* **316**: 1360–1368.

32. Hoffman, P.L., C.S. Rabe, F. Moses & B. Tabakoff. 1989. N-methyl-D-aspartate receptors and ethanol: inhibition of calcium flux and cyclic GMP production. *J. Neurochem.* **52**: 1937–1940.
33. Lima-Landman, M.T. & E.X. Albuquerque. 1989. Ethanol potentiates and blocks NMDA-activated single-channel currents in rat hippocampal pyramidal cells. *FEBS Lett.* **247**: 61–67.
34. Lovinger, D.M., G. White & F.F. Weight. 1989. Ethanol inhibits NMDA-activated ion current in hippocampal neurons. *Science* **243**: 1721–1724.
35. Wright, J.M., R.W. Peoples & F.F. Weight. 1996. Single-channel and whole-cell analysis of ethanol inhibition of NMDA-activated currents in cultured mouse cortical and hippocampal neurons. *Brain Res.* **738**: 249–256.
36. Nie, Z., S.G. Madamba & G.R. Siggins. 1994. Ethanol inhibits glutamatergic neurotransmission in nucleus accumbens neurons by multiple mechanisms. *J. Pharmacol. Exp. Ther.* **271**: 1566–1573.
37. Roberto, M., P. Schweitzer, S.G. Madamba, *et al.* 2004. Acute and chronic ethanol alter glutamatergic transmission in rat central amygdala: an *in vitro* and *in vivo* analysis. *J. Neurosci.* **24**: 1594–1603.
38. Ozawa, S., H. Kamiya & K. Tsuzuki. 1998. Glutamate receptors in the mammalian central nervous system. *Prog. Neurobiol.* **54**: 581–618.
39. Traynelis, S.F., L.P. Wollmuth, C.J. McBain, *et al.* 2010. Glutamate receptor ion channels: structure, regulation, and function. *Pharmacol. Rev.* **62**: 405–496.
40. Moriyoshi, K., M. Masu, T. Ishii, *et al.* 1991. Molecular cloning and characterization of the rat NMDA receptor. *Nature* **354**: 31–37.
41. Kuner, T., R. Schoepfer & E.R. Korpi. 1993. Ethanol inhibits glutamate-induced currents in heteromeric NMDA receptor subtypes. *NeuroReport* **5**: 297–300.
42. Lovinger, D.M. 1995. Developmental decrease in ethanol inhibition of N-methyl-D-aspartate receptors in rat neocortical neurons: relation to the actions of ifenprodil. *J. Pharmacol. Exp. Ther.* **274**: 164–172.
43. Masood, K., C. Wu, U. Brauneis & F.F. Weight. 1994. Differential ethanol sensitivity of recombinant N-methyl-D-aspartate receptor subunits. *Mol. Pharmacol.* **45**: 324–329.
44. Mirshahi, T. & J.J. Woodward. 1995. Ethanol sensitivity of heteromeric NMDA receptors: effects of subunit assembly, glycine and NMDAR1 Mg<sup>2+</sup>-insensitive mutants. *Neuropharmacology* **34**: 347–355.
45. Smothers, C.T., R. Clayton, T. Blevins & J.J. Woodward. 2001. Ethanol sensitivity of recombinant human N-methyl-D-aspartate receptors. *Neurochem. Int.* **38**: 333–340.
46. Chu, B., V. Anantharam & S.N. Treisman. 1995. Ethanol inhibition of recombinant heteromeric NMDA channels in the presence and absence of modulators. *J. Neurochem.* **65**: 140–148.
47. Wills, T.A., J.R. Klug, Y. Silberman, *et al.* 2012. GluN2B subunit deletion reveals key role in acute and chronic ethanol sensitivity of glutamate synapses in bed nucleus of the stria terminalis. *Proc. Natl. Acad. Sci. U.S.A.* **109**: E278–E287.
48. Suvarna, N., S.L. Borgland, J. Wang, *et al.* 2005. Ethanol alters trafficking and functional N-methyl-D-aspartate receptor NR2 subunit ratio via H-Ras. *J. Biol. Chem.* **280**: 31450–31459.
49. Brodie, M.S., S.A. Shefner & T.V. Dunwiddie. 1990. Ethanol increases the firing rate of dopamine neurons of the rat ventral tegmental area *in vitro*. *Brain Res.* **508**: 65–69.
50. Brodie, M.S., C. Pesold & S.B. Appel. 1999. Ethanol directly excites dopaminergic ventral tegmental area reward neurons. *Alcohol Clin. Exp. Res.* **23**: 1848–1852.
51. DiChiara, G. & A. Imperato. 1988. Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc. Natl. Acad. Sci. U.S.A.* **85**: 5274–5278.
52. Budygin, E.A., P.E. Phillips, D.L. Robinson, *et al.* 2001. Effect of acute ethanol on striatal dopamine neurotransmission in ambulatory rats. *J. Pharmacol. Exp. Ther.* **297**: 27–34.
53. Aryal, P., H. Dvir, S. Choe & P.A. Slesinger. 2009. A discrete alcohol pocket involved in GIRK channel activation. *Nat. Neurosci.* **12**: 988–995.
54. Beckstead, M.J. & T.J. Phillips. 2009. Mice selectively bred for high- or low-alcohol-induced locomotion exhibit differences in dopamine neuron function. *J. Pharmacol. Exp. Ther.* **329**: 342–349.
55. Ron, D. & R.O. Messing. 2013. Signaling pathways mediating alcohol effects. *Curr. Top. Behav. Neurosci.* **13**: 87–126.
56. Weiss, F., L.H. Parsons, G. Schulteis, *et al.* 1996. Ethanol self-administration restores withdrawal-associated deficiencies in accumbal dopamine and 5-hydroxytryptamine release in dependent rats. *J. Neurosci.* **16**: 3474–3485.
57. Yoshimoto, K., W.J. McBride, L. Lumeng & T.K. Li. 1992. Alcohol stimulates the release of dopamine and serotonin in the nucleus accumbens. *Alcohol* **9**: 17–22.
58. Yoshimoto, K., W.J. McBride, L. Lumeng & T.K. Li. 1992. Ethanol enhances the release of dopamine and serotonin in the nucleus accumbens of HAD and LAD lines of rats. *Alcohol Clin. Exp. Res.* **16**: 781–785.
59. Yoshimoto, K. & W.J. McBride. 1992. Regulation of nucleus accumbens dopamine release by the dorsal raphe nucleus in the rat. *Neurochem. Res.* **17**: 401–407.
60. Thielen, R.J., D.J. Bare, W.J. McBride, *et al.* 2002. Ethanol-stimulated serotonin release in the ventral hippocampus: an absence of rapid tolerance for the alcohol-preferring P rat and insensitivity in the alcohol-nonpreferring NP rat. *Pharmacol. Biochem. Behav.* **71**: 111–117.
61. Lovinger, D.M. 1991. Ethanol potentiation of 5-HT<sub>3</sub> receptor-mediated ion current in NCB-20 neuroblastoma-cells. *Neurosci. Lett.* **122**: 57–60.
62. Lovinger, D.M. & Q. Zhou. 1994. Alcohols potentiate ion current mediated by recombinant 5-HT<sub>3</sub>RA receptors expressed in a mammalian cell line. *Neuropharmacology* **33**: 1567–1572.
63. Sellers, E.M., G.A. Higgins & M.B. Sobell. 1992. 5-HT and alcohol abuse. *Trends Pharmacol. Sci.* **13**: 69.
64. Sellers, E.M., T. Toneatto, M.K. Romach, *et al.* 1994. Clinical efficacy of the 5-HT<sub>3</sub> antagonist ondansetron in alcohol abuse and dependence. *Alcohol Clin. Exp. Res.* **18**: 879–885.
65. Higley, J., M. Hasert, S. Suomi & M. Linnoila. 1998. The serotonin reuptake inhibitor sertraline reduces excessive alcohol consumption in nonhuman primates: effect of stress. *Neuropsychopharmacology* **18**: 431–443.

66. Heinz, A., J.D. Higley, J.G. Gorey, *et al.* 1998. In vivo association between alcohol intoxication, aggression, and serotonin transporter availability in nonhuman primates. *Am. J. Psychiatry* **155**: 1023–1028.
67. Lovinger, D.M. 1999. 5-HT<sub>3</sub> receptors and the neural actions of alcohols: an increasingly exciting topic. *Neurochem. Int.* **35**: 125–130.
68. Tikkanen, R., R.L. Sjöberg, F. Ducci, *et al.* 2009. Effects of MAOA-genotype, alcohol consumption, and aging on violent behavior. *Alcohol Clin. Exp. Res.* **33**: 428–434.
69. Maguire, E.P., E.A. Mitchell, S.J. Greig, *et al.* 2014. Extrasynaptic glycine receptors of rodent dorsal raphe serotonergic neurons: a sensitive target for ethanol. *Neuropsychopharmacology* **39**: 1232–1244.
70. Lowery-Gionta, E.G., C.A. Marcinkiewicz & T.L. Kash. 2015. Functional alterations in the dorsal raphe nucleus following acute and chronic ethanol exposure. *Neuropsychopharmacology* **40**: 590–600.
71. Vasudeva, R.K., A.R. Hobby & L.G. Kirby. 2015. Ethanol consumption in the Sprague–Dawley rat increases sensitivity of the dorsal raphe nucleus to 5,7-dihydroxytryptamine. *Behav. Brain. Res.* **S0166-4328**: 30044–30049.
72. Hwa, L.S., A. Shimamoto, T. Kayyali, *et al.* 2014. Dissociation of  $\mu$ -opioid receptor and CRF-R1 antagonist effects on escalated ethanol consumption and mPFC serotonin in C57BL/6J mice. *Addict. Biol.* doi: 10.1111/adb.12189.
73. McClintick, J.N., W.J. McBride, R.L. Bell, *et al.* 2015. Gene expression changes in serotonin, GABA<sub>A</sub> receptors, neuropeptides and ion channels in the dorsal raphe nucleus of adolescent alcohol-preferring (P) rats following binge-like alcohol drinking. *Pharmacol. Biochem. Behav.* **129**: 87–96.
74. Bach, H., V. Arango, S.A. Kassir, *et al.* 2014. Alcoholics have more tryptophan hydroxylase 2 mRNA and protein in the dorsal and median raphe nuclei. *Alcohol Clin. Exp. Res.* **38**: 1894–1901.
75. Miczek, K.A., J.F. DeBold, A.M.M. VanErp & W. Tornatzky. 1997. “Alcohol, GABA<sub>A</sub>-benzodiazepine receptor complex, and aggression.” In *Recent Developments in Alcoholism. Alcoholism and Violence*. M. Galanter, Ed.: 139–171. New York: Plenum.
76. Miczek, K.A. & R.M.M. deAlmeida. 2001. Oral drug self-administration in the home cage of mice: alcohol-heightened aggression and inhibition by the 5-HT<sub>1B</sub> agonist anpirtoline. *Psychopharmacology* **157**: 421–429.
77. Hwa, L.S., A.J. Nathanson, A. Shimamoto, *et al.* 2015. Aggression and increased glutamate in the mPFC during withdrawal from intermittent alcohol in outbred mice. *Psychopharmacology (Berl.)* **232**: 2889–2902.
78. Pohorecky, L.A. 1977. Biphasic action of ethanol. *Biobehav. Rev.* **1**: 231–240.
79. Cherek, D.R., J.L. Steinberg & B.R. Manno. 1985. Effects of alcohol on human aggressive behavior. *J. Stud. Alcohol* **46**: 321–328.
80. Miczek, K.A., E.M. Weerts, W. Tornatzky, *et al.* 1992. Alcohol and “bursts” of aggressive behavior: ethological analysis of individual differences in rats. *Psychopharmacology* **107**: 551–563.
81. Miczek, K.A., H.M. Barros, L. Sakoda & E.M. Weerts. 1998. Alcohol and heightened aggression in individual mice. *Alcohol Clin. Exp. Res.* **22**: 1698–1705.
82. Weerts, E.M., W. Tornatzky & K.A. Miczek. 1993. Prevention of the proaggressive effects of alcohol by benzodiazepine receptor antagonists in rats and in squirrel monkeys. *Psychopharmacology* **111**: 144–152.
83. Giancola, P.R. & A. Zeichner. 1997. The biphasic effects of alcohol on human physical aggression. *J. Abnorm. Psychol.* **106**: 598–607.
84. de Almeida, R.M.M., J.K. Rowlett, J.M. Cook, *et al.* 2004. GABA<sub>A</sub>/ $\alpha$ 1 receptor agonists and antagonists: effects on species-typical and heightened aggressive behavior after alcohol self-administration in mice. *Psychopharmacology* **172**: 255–263.
85. Haller, J. & M.R. Kruk. 2006. Normal and abnormal aggression: human disorders and novel laboratory models. *Neurosci. Biobehav. Rev.* **30**: 292–303.
86. Miczek, K.A., S.F. de Boer & J. Haller. 2013. Excessive aggression as model of violence: a critical evaluation of current preclinical methods. *Psychopharmacology* **226**: 445–458.
87. Peterson, J.T. & L.A. Pohorecky. 1989. Effect of chronic ethanol administration on intermale aggression in rats. *Aggress. Behav.* **15**: 201–215.
88. Citrome, L. & J. Volavka. 1999. Violent patients in the emergency setting. *Psychiatr. Clin. North Am.* **22**: 789–801.
89. Jaffe, J.H., T.F. Babor & D.H. Fishbein. 1988. Alcoholics, aggression and antisocial personality. *J. Stud. Alcohol* **1988**: 211–218.
90. Mayfield, D. 1976. Alcoholism, alcohol, intoxication and assaultive behavior. *Dis. Nerv. Syst.* **37**: 288–291.
91. Nicol, A.R., J.C. Gunn, J. Gristwood, *et al.* 1973. The relationship of alcoholism to violent behaviour resulting in long-term imprisonment. *Br. J. Psychiatry* **123**: 47–51.
92. Schuckit, M.A. & J.W. Russell. 1984. An evaluation of primary alcoholics with histories of violence. *J. Clin. Psychiatry* **45**: 3–6.
93. Miczek, K.A., S. Faccidomo, R.M.M. de Almeida, *et al.* 2004. Escalated aggressive behavior: new pharmacotherapeutic approaches and opportunities. *Ann. N. Y. Acad. Sci.* **1036**: 336–355.
94. Matsushita, S., M. Kimura, T. Miyakawa, *et al.* 2004. Association study of brain-derived neurotrophic factor gene polymorphism and alcoholism. *Alcohol Clin. Exp. Res.* **28**: 1609–1612.
95. Nilsson, K.W., R.L. Sjöberg, H.L. Wargelius, *et al.* 2007. The monoamine oxidase A (MAO-A) gene, family function and maltreatment as predictors of destructive behaviour during male adolescent alcohol consumption. *Addiction* **102**: 389–398.
96. Newman, E.L., A. Chu, B. Bahamón, *et al.* 2012. NMDA receptor antagonism: escalation of aggressive behavior in alcohol-drinking mice. *Psychopharmacology* **224**: 167–177.
97. Faccidomo, S., I.M. Quadros, A. Takahashi, *et al.* 2012. Infralimbic and dorsal raphe microinjection of the 5-HT<sub>1B</sub> receptor agonist CP-93129: attenuation of aggressive behavior in CFW male mice. *Psychopharmacology* **222**: 117–128.
98. Takahashi, A., C. Kwa, J.F. DeBold & K.A. Miczek. 2010. GABA<sub>A</sub> receptors in the dorsal raphe nucleus of mice: escalation of aggression after alcohol consumption. *Psychopharmacology* **211**: 467–477.

99. Chiavegatto, S., I.M. Quadros, G. Ambar & K.A. Miczek. 2010. Individual vulnerability to escalated aggressive behavior by a low dose of alcohol: decreased serotonin receptor mRNA in the prefrontal cortex of male mice. *Genes Brain Behav.* **9**: 110–119.
100. de Almeida, R.M., Q. Benini, J.S. Betat, *et al.* 2008. Heightened aggression after chronic flunitrazepam in male rats: potential links to cortical and caudate-putamen-binding sites. *Psychopharmacology* **197**: 309–318.
101. Fish, E.W., S.D. McKenzie-Quirk, M. Bannai & K.A. Miczek. 2008. 5-HT<sub>1B</sub> receptor inhibition of alcohol-heightened aggression in mice: comparison to drinking and running. *Psychopharmacology* **197**: 145–156.
102. de Almeida, R.M.M., E.M. Nikulina, S. Faccidomo, *et al.* 2001. Zolmitriptan—a 5-HT<sub>1B/D</sub> agonist, alcohol, and aggression in mice. *Psychopharmacology* **157**: 131–141.
103. Higley, J.D. 2001. Individual differences in alcohol-induced aggression. A nonhuman-primate model. *Alcohol Res. Health* **25**: 12–19.
104. Heinz, A.J., A. Beck, A. Meyer-Lindenberg, *et al.* 2011. Cognitive and neurobiological mechanisms of alcohol-related aggression. *Nat. Rev. Neurosci.* **12**: 400–413.
105. Miczek, K.A. & H. Barry III. 1977. Effects of alcohol on attack and defensive-submissive reactions in rats. *Psychopharmacology* **52**: 231–237.
106. Miczek, K.A., J.T. Winslow & J.F. DeBold. 1984. Heightened aggressive behavior by animals interacting with alcohol-treated conspecifics: studies with mice, rats and squirrel monkeys. *Pharmacol. Biochem. Behav.* **20**: 349–353.
107. Yoshimura, H. & N. Ogawa. 1983. Pharmacological analysis of agonistic behavior between resident and intruder mice: effects of ethylalcohol. *Nihon. Yakurigaku. Zasshi.* **81**: 135–141.
108. Sinha, R. 2001. How does stress increase risk of drug abuse and relapse? *Psychopharmacology* **158**: 343–359.
109. Brady, K.T. & S.C. Sonne. 1999. The role of stress in alcohol use, alcoholism treatment, and relapse. *Alcohol Res. Health* **23**: 263–271.
110. Pohorecky, L.A. 1981. The interaction of alcohol and stress, a review. *Neurosci. Biobehav. Rev.* **5**: 209–229.
111. Pohorecky, L.A. 1991. Stress and alcohol interaction: an update of human research. *Alcohol Clin. Exp. Res.* **15**: 438–459.
112. Becker, H.C., M.F. Lopez & T.L. Doremus-Fitzwater. 2011. Effects of stress on alcohol drinking: a review of animal studies. *Psychopharmacology* **218**: 131–156.
113. Noori, H.R., S. Helinski & R. Spanagel. 2014. Cluster and meta-analyses on factors influencing stress-induced alcohol drinking and relapse in rodents. *Addict. Biol.* **19**: 225–232.
114. Miczek, K.A., J.J. Yap & H.E. Covington III. 2008. Social stress, therapeutics and drug abuse: preclinical models of escalated and depressed intake. *Pharmacol. Ther.* **120**: 102–128.
115. McKenzie-Quirk, S.D. & K.A. Miczek. 2008. Social rank and social separation as determinants of alcohol drinking in squirrel monkeys. *Psychopharmacology* **201**: 137–145.
116. Helms, C.M., M.N. McClintick & K.A. Grant. 2012. Social rank, chronic ethanol self-administration, and diurnal pituitary-adrenal activity in cynomolgus monkeys. *Psychopharmacology* **224**: 133–143.
117. Kudryavtseva, N.N., I.V. Bakshtanovskaya & L.A. Koryakina. 1991. Social model of depression in mice of C57BL/6J strain. *Pharmacol. Biochem. Behav.* **38**: 315–320.
118. Peters, S., D.A. Slattery, P.J. Flor, *et al.* 2013. Differential effects of baclofen and oxytocin on the increased ethanol consumption following chronic psychosocial stress in mice. *Addict. Biol.* **18**: 66–77.
119. Sillaber, I., G. Rammes, S. Zimmermann, *et al.* 2002. Enhanced and delayed stress-induced alcohol drinking in mice lacking functional CRH1 receptors. *Science* **296**: 931–933.
120. Norman, K.J., J.A. Seiden, J.A. Klickstein, *et al.* 2015. Social stress and escalated drug self-administration in mice I. Alcohol and corticosterone. *Psychopharmacology* **232**: 991–1001.
121. Molander, A., V. Vengeliene, M. Heilig, *et al.* 2012. Brain-specific inactivation of the Crhr1 gene inhibits post-dependent and stress-induced alcohol intake, but does not affect relapse-like drinking. *Neuropsychopharmacology* **37**: 1047–1056.
122. Riga, D., L.J. Schmitz, J.E. Van der Harst, *et al.* 2014. A sustained depressive state promotes a guanfacine reversible susceptibility to alcohol seeking in rats. *Neuropsychopharmacology* **39**: 1115–1124.
123. Scott, J.P. 1958. *Aggression*. Chicago: University of Chicago Press.
124. Fish, E.W., J.F. DeBold & K.A. Miczek. 2002. Aggressive behavior as a reinforcer in mice: activation by allopregnanolone. *Psychopharmacology* **163**: 459–466.
125. Hsu, Y., R.L. Earley & L.L. Wolf. 2006. Modulation of aggressive behaviour by fighting experience: mechanisms and contest outcomes. *Biol. Rev. Camb. Philos. Soc.* **81**: 33–74.
126. Nelson, R.J. & B.C. Trainor. 2007. Neural mechanisms of aggression. *Nat. Neurosci. Rev.* **8**: 536–546.
127. Caramaschi, D., S.F. de Boer, H. de Vries & J.M. Koolhaas. 2008. Development of violence in mice through repeated victory along with changes in prefrontal cortex neurochemistry. *Behav. Brain Res.* **189**: 263–272.
128. Couppis, M.H. & C.H. Kennedy. 2008. The rewarding effect of aggression is reduced by nucleus accumbens dopamine receptor antagonism in mice. *Psychopharmacology* **197**: 449–456.
129. Rassnick, S., L. Pulvirenti & G.F. Koob. 1992. Oral ethanol self-administration in rats is reduced by the administration of dopamine and glutamate receptor antagonists into the nucleus accumbens. *Psychopharmacology* **109**: 92–98.
130. Weiss, F., M.T. Lorang, F.E. Bloom & G.F. Koob. 1993. Oral alcohol self-administration stimulates dopamine release in the rat nucleus accumbens—genetic and motivational determinants. *J. Pharmacol. Exp. Ther.* **267**: 250–258.
131. Koob, G.F., A.J. Roberts, G. Schulteis, *et al.* 1998. Neurocircuitry targets in ethanol reward and dependence. *Alcohol Clin. Exp. Res.* **22**: 3–9.
132. Heilig, M. & M. Egli. 2006. Pharmacological treatment of alcohol dependence: target symptoms and target mechanisms. *Pharmacol. Ther.* **111**: 855–876.

133. Barr, C.S., T.K. Newman, S. Lindell, *et al.* 2004. Early experience and sex interact to influence limbic–hypothalamic–pituitary–adrenal-axis function after acute alcohol administration in rhesus macaques (*Macaca mulatta*). *Alcohol Clin. Exp. Res.* **28**: 1114–1119.
134. Rodd, Z.A., R.L. Bell, R.I. Melendez, *et al.* 2004. Comparison of intracranial self-administration of ethanol within the posterior ventral tegmental area between alcohol-preferring and Wistar rats. *Alcohol Clin. Exp. Res.* **28**: 1212–1219.
135. Spanagel, R. & F. Weiss. 1999. The dopamine hypothesis of reward: past and current status. *Trends Neurosci.* **22**: 521–527.
136. Heilig, M. & G.F. Koob. 2007. A key role for corticotropin-releasing factor in alcohol dependence. *Trends Neurosci.* **30**: 399–406.
137. Fish, E.W., J.F. De Bold & K.A. Miczek. 2005. Escalated aggression as a reward: corticosterone and GABA<sub>A</sub> receptor positive modulators in mice. *Psychopharmacology* **182**: 116–127.
138. van Erp, A.M. & K.A. Miczek. 2007. Increased accumbal dopamine during daily alcohol consumption and subsequent aggressive behavior in rats. *Psychopharmacology* **191**: 679–688.
139. de Boer, S.F., D. Caramaschi, D. Natarajan & J.M. Koolhaas. 2009. The vicious cycle towards violence: focus on the negative feedback mechanisms of brain serotonin neurotransmission. *Front Behav. Neurosci.* **3**: 52.
140. de Boer, S.F. & J.M. Koolhaas. 2005. 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptor agonists and aggression: a pharmacological challenge of the serotonin deficiency hypothesis. *Eur. J. Pharmacol.* **526**: 125–139.
141. Van Der Vegt, B.J., N. Lieuwes, E.H. van de Wall, *et al.* 2003. Activation of serotonergic neurotransmission during the performance of aggressive behavior in rats. *Behav. Neurosci.* **117**: 667–674.
142. Kravitz, E.A. & R. Huber. 2003. Aggression in invertebrates. *Curr. Opin. Neurobiol.* **13**: 736–743.
143. Summers, C.H., W.J. Korzan, J.L. Lukkes, *et al.* 2005. Does serotonin influence aggression? Comparing regional activity before and during social interaction. *Physiol. Biochem. Zool.* **78**: 679–694.
144. Miczek, K.A., R.M.M. de Almeida, E.A. Kravitz, *et al.* 2007. Neurobiology of escalated aggression and violence. *J. Neurosci.* **27**: 11803–11806.
145. Virkkunen, M. & M. Linnoila. 1993. Brain serotonin, type II alcoholism and impulsive violence. *J. Stud. Alcohol Suppl.* **11**: 163–169.
146. Virkkunen, M., E. Kallio, R. Rawlings, *et al.* 1994. Personality profiles and state aggressiveness in Finnish alcoholic, violent offenders, fire setters, and healthy volunteers. *Arch. Gen. Psychiatry* **51**: 28–33.
147. Virkkunen, M., D. Goldman, D.A. Nielsen & M. Linnoila. 1995. Low brain serotonin turnover rate (low CSF 5-HIAA) and impulsive violence. *J. Psychiatry Neurosci.* **20**: 271–275.
148. Beck, A., A.J. Heinz & A. Heinz. 2014. Translational clinical neuroscience perspectives on the cognitive and neurobiological mechanisms underlying alcohol-related aggression. *Curr. Top. Behav. Neurosci.* **17**: 443–474.
149. Brodie, B.B. & P.A. Shore. 1957. A concept for a role of serotonin and norepinephrine as chemical mediators in the brain. *Analgesia* **66**: 631–642.
150. Giacalone, E., M. Tansella, L. Valzelli & S. Garattini. 1968. Brain serotonin metabolism in isolated aggressive mice. *Biochem. Pharmacol.* **17**: 1315–1327.
151. Olivier, B. 2004. Serotonin and aggression. *Ann. N. Y. Acad. Sci.* **1036**: 382–392.
152. Takahashi, A., I.M. Quadros, R.M. de Almeida & K.A. Miczek. 2011. Brain serotonin receptors and transporters: initiation vs. termination of escalated aggression. *Psychopharmacology* **213**: 183–212.
153. Morrison, T.R. & R.H. Melloni, Jr. 2014. The role of serotonin, vasopressin, and serotonin/vasopressin interactions in aggressive behavior. *Curr. Top. Behav. Neurosci.* **17**: 189–228.
154. Lesch, K.P., D. Bengel, A. Heils, *et al.* 1996. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* **264**: 1537–1551.
155. Schwandt, M.L., S.G. Lindell, R.L. Sjöberg, *et al.* 2010. Gene–environment interactions and response to social intrusion in male and female rhesus macaques. *Biol. Psychiatry* **67**: 323–330.
156. de Almeida, R.M.M., P.F. Ferrari, S. Parmigiani & K.A. Miczek. 2005. Escalated aggressive behavior: dopamine, serotonin and GABA. *Eur. J. Pharmacol.* **526**: 51–64.
157. de Almeida, R.M., M.M. Rosa, D.M. Santos, *et al.* 2006. 5-HT<sub>1B</sub> receptors, ventral orbitofrontal cortex, and aggressive behavior in mice. *Psychopharmacology* **185**: 441–450.
158. de Almeida, R.M.M. & K.A. Miczek. 2002. Aggression escalated by social instigation or by discontinuation of reinforcement (“frustration”) in mice: inhibition by anpirtoline, a 5-HT<sub>1B</sub> receptor agonist. *Neuropsychopharmacology* **27**: 171–181.
159. Gowin, J.L., A.C. Swann, F.G. Moeller & S.D. Lane. 2010. Zolmitriptan and human aggression: interaction with alcohol. *Psychopharmacology* **210**: 521–531.
160. Coccaro, E.F., M.S. McCloskey, D.A. Fitzgerald & K.L. Phan. 2007. Amygdala and orbitofrontal reactivity to social threat in individuals with impulsive aggression. *Biol. Psychiatry* **62**: 168–178.
161. Sterzer, P., C. Stadler, A. Krebs, *et al.* 2005. Abnormal neural responses to emotional visual stimuli in adolescents with conduct disorder. *Biol. Psychiatry* **57**: 7–15.
162. Lee, T.M., S.C. Chan & A. Raine. 2008. Strong limbic and weak frontal activation to aggressive stimuli in spouse abusers. *Mol. Psychiatry* **13**: 655–656.
163. Kollack-Walker, S. & S.W. Newman. 1995. Mating and agonistic behavior produce different patterns of Fos immunolabeling in the male Syrian hamster brain. *Neuroscience* **66**: 721–736.
164. Lee, H., D.W. Kim, R. Remedios, *et al.* 2014. Scalable control of mounting and attack by Esr1+ neurons in the ventromedial hypothalamus. *Nature* **509**: 627–632.
165. Lin, D., M.P. Boyle, P. Dollar, *et al.* 2011. Functional identification of an aggression locus in the mouse hypothalamus. *Nature* **470**: 221–226.
166. Bannai, M., E.W. Fish, S. Faccidomo & K.A. Miczek. 2007. Anti-aggressive effects of agonists at 5-HT<sub>1B</sub> receptors in

- the dorsal raphe nucleus of mice. *Psychopharmacology* **193**: 295–304.
167. Faccidomo, S., M. Bannai & K.A. Miczek. 2008. Escalated aggression after alcohol drinking in male mice: dorsal raphe and prefrontal cortex serotonin and 5-HT<sub>1B</sub> receptors. *Neuropsychopharmacology* **33**: 2888–2899.
  168. Mhatre, M.C., G. Pena, W. Sieghart & M.K. Ticku. 1993. Antibodies specific for GABA<sub>A</sub> receptor  $\alpha$  subunits reveal that chronic alcohol treatment down-regulates  $\alpha$ -subunit expression in rat brain regions. *J. Neurochem.* **61**: 1620–1625.
  169. Matthews, D.B., L.L. Devaud, J.M. Fritschy, *et al.* 1998. Differential regulation of GABA<sub>A</sub> receptor gene expression by ethanol in the rat hippocampus versus cerebral cortex. *J. Neurochem.* **70**: 1160–1166.
  170. Follesa, P., F. Biggio, G. Talani, *et al.* 2006. Neurosteroids, GABA<sub>A</sub> receptors, and ethanol dependence. *Psychopharmacology* **186**: 267–280.
  171. Mahmoudi, M., M.H. Kang, N. Tillakaratne, *et al.* 1997. Chronic intermittent ethanol treatment in rats increases GABA<sub>A</sub> receptor  $\alpha 4$ -subunit expression: possible relevance to alcohol dependence. *J. Neurochem.* **68**: 2485–2492.
  172. Miczek, K.A., J.F. DeBold & A.M.M. Van Erp. 1994. Neuropharmacological characteristics of individual differences in alcohol effects on aggression in rodents and primates. *Behav. Pharmacol.* **5**: 407–421.
  173. DiMascio, A. 1973. The effects of benzodiazepines on aggression: reduced or increased? *Psychopharmacologia* **30**: 95–102.
  174. Albrecht, B., P.K. Staiger, K. Hall, *et al.* 2014. Benzodiazepine use and aggressive behaviour: a systematic review. *Aust. N. Z. J. Psychiatry* **48**: 1096–1114.
  175. Jonas, J.M., B.S. Coleman, A.Q. Sheridan & R.W. Kalinske. 1992. Comparative clinical profiles of triazolam versus other shorter-acting hypnotics. *J. Clin. Psychiatry* **53**(Suppl.): 19–31.
  176. Pilowsky, L.S., H. Ring, P.J. Shine, *et al.* 1992. Rapid tranquillisation. A survey of emergency prescribing in a general psychiatric hospital. *Br. J. Psychiatry* **160**: 831–835.
  177. Fulde, G. & P. Preisz. 2011. Managing aggressive and violent patients. *Aust. Prescr.* **34**: 115–118.
  178. Angelotti, T.P. & R.L. Macdonald. 1993. Assembly of GABA<sub>A</sub> receptor subunits:  $\alpha 1$ - $\beta 1$  and  $\alpha 1$ - $\beta 1$ - $\gamma 2S$  subunits produce unique ion channels with dissimilar single-channel properties. *J. Neurosci.* **13**: 1429–1440.
  179. Pritchett, D.B., H. Sontheimer, B.D. Shivers, *et al.* 1989. Importance of a novel GABA<sub>A</sub> receptor subunit for benzodiazepine pharmacology. *Nature* **338**: 582–585.
  180. Günther, U., J. Benson, D. Benke, *et al.* 1995. Benzodiazepine-insensitive mice generated by targeted disruption of the  $\gamma 2$  subunit gene of  $\gamma$ -aminobutyric acid type A receptors. *Proc. Natl. Acad. Sci. U.S.A.* **92**: 7749–7753.
  181. Tan, K.R., U. Rudolph & C. Luscher. 2011. Hooked on benzodiazepines: GABA<sub>A</sub> receptor subtypes and addiction. *Trends Neurosci.* **34**: 188–197.
  182. Newman, E.L., K.S. Smith, A. Takahashi, *et al.* 2015.  $\alpha 2$ -containing GABA(A) receptors: a requirement for midazolam-escalated aggression and social approach in mice. *Psychopharmacology*. In press.
  183. Puglisi-Allegra, S. & P. Mandel. 1980. Effects of sodium n-dipropylacetate, muscimol hydrobromide and (R,S)nipecotinic acid amide on isolation-induced aggressive behavior in mice. *Psychopharmacology* **70**: 287–290.
  184. Puglisi-Allegra, S., S. Simler, E. Kempf & P. Mandel. 1981. Involvement of the GABAergic system on shock-induced aggressive behavior in two strains of mice. *Pharmacol. Biochem. Behav.* **14**(Suppl. 1): 13–18.
  185. Rodgers, R.J. & A. Depaulis. 1982. Gabaergic influences on defensive fighting in rats. *Pharmacol. Biochem. Behav.* **17**: 451–456.
  186. Allikmets, L.H. & L.K. Rago. 1983. The action of benzodiazepine antagonist Ro 15–1788 on the effects of GABAergic drugs. *Naunyn. Schmiedebergs. Arch. Pharmacol.* **324**: 235–237.
  187. Takahashi, A., A. Shimamoto, C.O. Boyson, *et al.* 2010. GABA<sub>B</sub> receptor modulation of serotonin neurons in the dorsal raphe nucleus and escalation of aggression in mice. *J. Neurosci.* **30**: 11771–11780.
  188. Majewska, M.D., N.L. Harrison, R.D. Schwartz, *et al.* 1986. Steroid hormone metabolites are barbiturate-like modulators of the GABA receptor. *Science* **232**: 1004–1007.
  189. Gee, K.W., W.C. Chang, R.E. Brinton & B.S. McEwen. 1987. GABA-dependent modulation of the Cl<sup>-</sup> ionophore by steroids in rat brain. *Eur. J. Pharmacol.* **136**: 419–423.
  190. Baulieu, E.E. & P. Robel. 1990. Neurosteroids: a new brain function? *J. Steroid Biochem. Mol. Biol.* **37**: 395–403.
  191. Akwa, Y., J. Young, K. Kabbadj, *et al.* 1991. Neurosteroids: biosynthesis, metabolism and function of pregnenolone and dehydroepiandrosterone in the brain. *J. Steroid Biochem. Mol. Biol.* **40**: 71–81.
  192. Baulieu, E.E. 1992. Neurosteroids: an overview. *Adv. Biochem. Psychopharmacol.* **47**: 1–16.
  193. Lambert, J.L., D. Belilli, C. Hill-Venning & J.A. Peters. 1995. Neurosteroids and GABA<sub>A</sub> receptor function. *Trends Pharmacol. Sci.* **16**: 295–303.
  194. Schüle, C., C. Nothdurfter & R. Rupprecht. 2014. The role of allopregnanolone in depression and anxiety. *Prog. Neurobiol.* **113**: 79–87.
  195. VanDoren, M.J., D.B. Matthews, G.C. Janis, *et al.* 2000. Neuroactive steroid 3 $\alpha$ -hydroxy-5 $\alpha$ -pregnane-20-one modulates electrophysiological and behavioral actions of ethanol. *J. Neurosci.* **20**: 1982–1989.
  196. Morrow, A.L., G.C. Janis, M.J. VanDoren, *et al.* 1999. Neurosteroids mediate pharmacological effects of ethanol: a new mechanism of ethanol action? *Alcohol. Clin. Exp. Res.* **23**: 1933–1940.
  197. Purdy, R.H., A.L. Morrow, P.H. Moore & S.M. Paul. 1991. Stress-induced elevations of  $\gamma$ -aminobutyric acid type A receptor-active steroids in the rat brain. *Proc. Natl. Acad. Sci. U.S.A.* **88**: 4553–4557.
  198. Pinna, G., E. Costa & A. Guidotti. 2005. Changes in brain testosterone and allopregnanolone biosynthesis elicit aggressive behavior. *Proc. Natl. Acad. Sci. U.S.A.* **102**: 2135–2140.
  199. Nelson, M. & G. Pinna. 2011. S-norfluoxetine microinfused into the basolateral amygdala increases allopregnanolone levels and reduces aggression in socially isolated mice. *Neuropharmacology* **60**: 1154–1159.

200. Fish, E.W., S. Faccidomo, J.F. DeBold & K.A. Miczek. 2001. Alcohol, allopregnanolone and aggression in mice. *Psychopharmacology* **153**: 473–483.
201. Guidotti, A., E. Dong, K. Matsumoto, *et al.* 2001. The socially-isolated mouse: a model to study the putative role of allopregnanolone and 5 $\alpha$ -dihydroprogesterone in psychiatric disorders. *Brain Res. Rev.* **37**: 110–115.
202. Biggio, G., M.G. Pisu, F. Biggio & M. Serra. 2014. Allopregnanolone modulation of HPA axis function in the adult rat. *Psychopharmacology* **231**: 3437–3444.
203. Addolorato, G., L. Leggio, L. Abenavoli, *et al.* 2006. Baclofen in the treatment of alcohol withdrawal syndrome: a comparative study vs diazepam. *Am. J. Med.* **119**: 276–278.
204. Brennan, J.L., J.G. Leung, J.P. Gagliardi, *et al.* 2013. Clinical effectiveness of baclofen for the treatment of alcohol dependence: a review. *Clin. Pharmacol.* **5**: 99–107.
205. Ponzovsky, A.M., P. Rosca, E. Aronovich, *et al.* 2014. Baclofen as add-on to standard psychosocial treatment for alcohol dependence: a randomized, double-blind, placebo-controlled trial with 1-year follow-up. *J. Subst. Abuse Treat.* **52**: 24–30.
206. Bucknam, W. 2007. Suppression of symptoms of alcohol dependence and craving using high-dose baclofen. *Alcohol Alcohol.* **42**: 158–160.
207. Addolorato, G., F. Caputo, E. Capristo, *et al.* 2002. Baclofen efficacy in reducing alcohol craving and intake: a preliminary double-blind randomized controlled study. *Alcohol Alcohol.* **37**: 504–508.
208. File, S.E., A. Zharkovsky & K. Gulati. 1991. Effects of baclofen and nitrendipine on ethanol withdrawal responses in the rat. *Neuropharmacology* **30**: 183–190.
209. Rudissaar, R., K. Pruus, T. Skrebuhova-Malmros, *et al.* 2000. Involvement of GABAergic neurotransmission in the neurobiology of the apomorphine-induced aggressive behavior paradigm, a model of psychotic behavior in rats. *Methods Find Exp. Clin. Pharmacol.* **22**: 637–640.
210. Takahashi, A., A.N. Schilit, J. Kim, *et al.* 2012. Behavioral characterization of escalated aggression induced by GABA<sub>B</sub> receptor activation in the dorsal raphe nucleus. *Psychopharmacology* **224**: 155–166.
211. Gruol, D.L., K.L. Parsons & N. DiJulio. 1997. Acute ethanol alters calcium signals elicited by glutamate receptor agonists and K<sup>+</sup> depolarization in cultured cerebellar Purkinje neurons. *Brain Res.* **773**: 82–89.
212. Hassanain, M., S. Bhatt & A. Siegel. 2003. Differential modulation of feline defensive rage behavior in the medial hypothalamus by 5-HT<sub>1A</sub> and 5-HT<sub>2</sub> receptors. *Brain Res.* **981**: 201–209.
213. Fernandes, C., J.L. Paya-Cano, F. Sluyter, *et al.* 2004. Hippocampal gene expression profiling across eight mouse inbred strains: towards understanding the molecular basis for behaviour. *Eur. J. Neurosci.* **19**: 2576–2582.
214. Schubert, K., M.B. Shaikh, Y. Han, *et al.* 1996. Differential effects of ethanol on feline rage and predatory attack behavior: an underlying neural mechanism. *Alcohol. Clin. Exp. Res.* **20**: 882–889.
215. Enoch, M.A., A.A. Rosser, Z. Zhou, *et al.* 2014. Expression of glutamatergic genes in healthy humans across 16 brain regions; altered expression in the hippocampus after chronic exposure to alcohol or cocaine. *Genes Brain Behav.* **13**: 758–768.
216. Jin, Z., A.K. Bhandage, I. Bazov, *et al.* 2014. Selective increases of AMPA, NMDA, and kainate receptor subunit mRNAs in the hippocampus and orbitofrontal cortex but not in prefrontal cortex of human alcoholics. *Front. Cell Neurosci.* **8**: 11.
217. Cacciaglia, R., F. Nees, S.T. Pohlack, *et al.* 2013. A risk variant for alcoholism in the NMDA receptor affects amygdala activity during fear conditioning in humans. *Biol. Psychol.* **94**: 74–81.
218. Domart, M.C., A. Benyamina, A. Lemoine, *et al.* 2012. Association between a polymorphism in the promoter of a glutamate receptor subunit gene (GRIN2A) and alcoholism. *Addict. Biol.* **17**: 783–785.
219. Karpayak, V.M., J.M. Biernacka, J.R. Geske, *et al.* 2014. Genetic markers associated with abstinence length in alcohol-dependent subjects treated with acamprosate. *Transl. Psychiatry* **4**: e462.
220. Du, X., A.J. Elberger, D.B. Matthews & K.M. Hamre. 2012. Heterozygous deletion of NR1 subunit of the NMDA receptor alters ethanol-related behaviors and regional expression of NR2 subunits in the brain. *Neurotoxicol. Teratol.* **34**: 177–186.
221. Duncan, G.E., S.S. Moy, A. Perez, *et al.* 2004. Deficits in sensorimotor gating and tests of social behavior in a genetic model of reduced NMDA receptor function. *Behav. Brain Res.* **153**: 507–519.
222. Rammes, G., B. Mahal, J. Putzke, *et al.* 2001. The anti-craving compound acamprosate acts as a weak NMDA-receptor antagonist, but modulates NMDA-receptor subunit expression similar to memantine and MK-801. *Neuropharmacology* **40**: 749–760.
223. Hölter, S.M., A.C. Linthorst, J.M. Reul & R. Spanagel. 2000. Withdrawal symptoms in a long-term model of voluntary alcohol drinking in Wistar rats. *Pharmacol. Biochem. Behav.* **66**: 143–151.
224. Morales, M., E.I. Varlinskaya & L.P. Spear. 2013. Low doses of the NMDA receptor antagonists, MK-801, PEAQX, and ifenprodil, induces social facilitation in adolescent male rats. *Behav. Brain Res.* **250**: 18–22.
225. Krystal, J.H., I.L. Petrakis, D. Limoncelli, *et al.* 2003. Altered NMDA glutamate receptor antagonist response in recovering ethanol-dependent patients. *Neuropsychopharmacology* **28**: 2020–2028.
226. Krupitsky, E.M., O. Neznanova, D. Masalov, *et al.* 2007. Effect of memantine on cue-induced alcohol craving in recovering alcohol-dependent patients. *Am. J. Psychiatry* **164**: 519–523.
227. Krystal, J.H., J. Staley, G. Mason, *et al.* 2006. Gamma-aminobutyric acid type A receptors and alcoholism: intoxication, dependence, vulnerability, and treatment. *Arch. Gen. Psychiatry* **63**: 957–968.
228. Allgaier, C. 2002. Ethanol sensitivity of NMDA receptors. *Neurochem. Int.* **41**: 377–382.
229. Wenzel, A., L. Scheurer, R. Künzi, *et al.* 1995. Distribution of NMDA receptor subunit proteins NR2A, 2B, 2C and 2D in rat brain. *NeuroReport* **7**: 45–48.

230. Halasz, J., M. Toth, I. Kallo, *et al.* 2006. The activation of prefrontal cortical neurons in aggression—a double labeling study. *Behav. Brain Res.* **175**: 166–175.
231. Takahashi, A., K. Nagayasu, N. Nishitani, *et al.* 2014. Control of intermale aggression by medial prefrontal cortex activation in the mouse. *PLoS One* **9**: e94657.
232. Lau, C.G. & R.S. Zukin. 2007. NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat. Rev. Neurosci.* **8**: 413–426.
233. Ciccocioppo, R., D.R. Gehlert, A. Ryabinin, *et al.* 2009. Stress-related neuropeptides and alcoholism: CRH, NPY, and beyond. *Alcohol* **43**: 491–498.
234. Heinrichs, M., D.B. von & G. Domes. 2009. Oxytocin, vasopressin, and human social behavior. *Front. Neuroendocrinol.* **30**: 548–557.
235. Bielsky, I.F. & L.J. Young. 2004. Oxytocin, vasopressin, and social recognition in mammals. *Peptides* **25**: 1565–1574.
236. De Felipe, C., J.F. Herrero, J.A. O'Brien, *et al.* 1998. Altered nociception, analgesia and aggression in mice lacking the receptor for substance P. *Nature* **392**: 394–397.
237. Dunn, A.J. & C.W. Berridge. 1990. Physiological and behavioral responses to corticotropin-releasing factor administration: is CRF a mediator of anxiety or stress responses? *Brain Res. Rev.* **15**: 71–100.
238. Vale, W., J. Spiess, C. Rivier & J. Rivier. 1981. Characterization of a 41-residue ovine hypothalamic peptide that stimulates secretion of corticotropin and  $\beta$ -endorphin. *Science* **213**: 1394–1397.
239. Lowry, C.A. & F.L. Moore. 2006. Regulation of behavioral responses by corticotropin-releasing factor. *Gen. Comp. Endocrinol.* **146**: 19–27.
240. Hostetler, C.M. & A.E. Ryabinin. 2013. The CRF system and social behavior: a review. *Front. Neurosci.* **7**: 92.
241. Mele, A., S. Cabib, A. Oliverio, *et al.* 1987. Effects of corticotropin releasing factor and sauvagine on social behavior of isolated mice. *Peptides* **8**: 935–938.
242. Carpenter, R.E., W.J. Korzan, C. Bockholt, *et al.* 2009. Corticotropin releasing factor influences aggression and monoamines: modulation of attacks and retreats. *Neuroscience* **158**: 412–425.
243. Elkabir, D.R., M.E. Wyatt, S.V. Vellucci & J. Herbert. 1990. The effects of separate or combined infusions of corticotropin-releasing factor and vasopressin either intraventricularly or into the amygdala on aggressive and investigative behaviour in the rat. *Regul. Pept.* **28**: 199–214.
244. Farrokhi, C., D.C. Blanchard, G. Griebel, *et al.* 2004. Effects of the CRF1 antagonist SSR125543A on aggressive behaviors in hamsters. *Pharmacol. Biochem. Behav.* **77**: 465–469.
245. Cheng, S.Y. & Y. Delville. 2010. Play fighting and corticotropin-releasing hormone in the lateral septum of golden hamsters. *Neuroscience* **169**: 236–245.
246. Backström, T., A. Pettersson, V. Johansson & S. Winberg. 2011. CRF and urotensin I effects on aggression and anxiety-like behavior in rainbow trout. *J. Exp. Biol.* **214**: 907–914.
247. Quadros, I.M., L.S. Hwa, A. Shimamoto, *et al.* 2014. Prevention of alcohol-heightened aggression by CRF-R1 antagonists in mice: critical role for DRN-PFC serotonin pathway. *Neuropsychopharmacology* **39**: 2874–2883.
248. Anderson, B.M., M.C. Stevens, S.A. Meda, *et al.* 2011. Functional imaging of cognitive control during acute alcohol intoxication. *Alcohol Clin. Exp. Res.* **35**: 156–165.
249. Virkkunen, M. 1985. Urinary free cortisol secretion in habitually violent offenders. *Acta Psychiatr. Scand.* **72**: 40–44.
250. Hibbeln, J.R., G. Bissette, J.C. Umhau & D.T. George. 2004. Omega-3 status and cerebrospinal fluid corticotrophin releasing hormone in perpetrators of domestic violence. *Biol. Psychiatry* **56**: 895–897.
251. Lutz, P.E. & B.L. Kieffer. 2013. The multiple facets of opioid receptor function: implications for addiction. *Curr. Opin. Neurobiol.* **23**: 473–479.
252. MacHin, A.J. & R.I.M. Dunbar. 2011. The brain opioid theory of social attachment: a review of the evidence. *Behaviour* **148**: 985–1025.
253. Braud, W.G. & J.E. Weibel. 1969. Acquired stimulus control of drug-induced changes in aggressive display in *Betta splendens*. *J. Exp. Anal. Behav.* **12**: 773–777.
254. Avis, H.H. & H.V.S. Peeke. 1975. Differentiation by morphine of two types of aggressive behavior in the convict cichlid (*Cichlasoma nigrofasciatum*). *Psychopharmacologia* **43**: 287–288.
255. Faisal, M., F. Chiappelli, I.I. Ahmed, *et al.* 1989. Social confrontation “stress” in aggressive fish is associated with an endogenous opioid-mediated suppression of proliferative response to mitogens and nonspecific cytotoxicity. *Brain Behav. Immun.* **3**: 223–233.
256. Wikler, A. 1944. Studies on the action of morphine on the central nervous system of cat. *J. Pharmacol. Exp. Ther.* **80**: 176–187.
257. Cronin, G.M., P.R. Wiekema & J.M. van Ree. 1985. Endogenous opioids are involved in abnormal stereotyped behaviours of tethered sows. *Neuropeptides* **6**: 527–530.
258. Dyakonova, V., F.W. Schurmann & D.A. Sakharov. 2002. Effects of opiate ligands on intraspecific aggression in crickets. *Peptides* **23**: 835–841.
259. Espert, R., J.F. Navarro, A. Salvador & V.M. Simon. 1993. Effects of morphine hydrochloride on social encounters between male mice. *Aggress. Behav.* **19**: 377–383.
260. Winslow, J.T. & K.A. Miczek. 1988. Naltrexone blocks amphetamine-induced hyperactivity, but not disruption of social and agonistic behavior in mice and squirrel monkeys. *Psychopharmacology* **96**: 493–499.
261. Rodgers, R.J. & C.A. Hendrie. 1983. Social conflict activates status-dependent endogenous analgesic or hyperalgesic mechanisms in male mice: effects of naloxone on nociception and behaviour. *Physiol. Behav.* **30**: 775–780.
262. Dyakonova, V., F.W. Schormann & D.A. Sakharov. 2000. Social aggressiveness of female and subordinate male crickets is released by opiate receptor antagonist. *Acta Biol. Hung.* **51**: 363–367.
263. Borg, S., H. Kvande, U. Rydberg, *et al.* 1982. Endorphin levels in human cerebrospinal fluid during alcohol intoxication and withdrawal. *Psychopharmacology* **78**: 101–103.
264. Miczek, K.A., M.L. Thompson & L. Shuster. 1982. Opioid-like analgesia in defeated mice. *Science* **215**: 1520–1522.
265. Siegfried, B. & H.R. Frischknecht. 1989. Place avoidance learning and stress-induced analgesia in the attacked

- mouse: role of endogenous opioids. *Behav. Neural Biol.* **52**: 95–107.
266. Konicki, P.E. & S.C. Schulz. 1989. Rationale for clinical trials of opiate antagonists in treating patients with personality disorders and self-injurious behavior. *Psychopharmacol. Bull.* **25**: 556–563.
267. Dunwiddie, T. 2001. “Effects of ethanol on neurotransmitter systems.” In *Alcohol and Alcohol Actions Lecture Series*. D. Gruol & Y. Israel, Eds. Austin: Research Society on Alcoholism.
268. Meyer, H. 1899. Zur Theorie der Alkoholnarkose. I. Welche Eigenschaft der Anästhetica bedingt ihre narkotische Wirkung? *Naunyn. Schmiedebergs. Arch. Pharmacol. Exp. Pathol.* **42**: 109–118.
269. Meyer, H. 1901. Zur Theorie der Alkoholnarkose. III. Der Einfluss wechselnder Temperatur auf Wirkungsstärke und Theilungscoefficient der Narcoticanarkose. *Naunyn. Schmiedebergs. Arch. Pharmacol. Exp. Pathol.* **46**: 338–346.
270. Overton, E. 1901. *Studien über die Narkose: zugleich ein Beitrag zur allgemeinen Pharmakologie*. Jena: Gustav Fischer.
271. Lovinger, D.M. 1997. Alcohols and neurotransmitter gated ion channels: past, present and future. *Naunyn. Schmiedebergs. Arch. Pharmacol.* **356**: 267–282.
272. Franks, N.P. & W.R. Lieb. 1984. Do general anaesthetics act by competitive binding to specific receptors? *Nature* **310**: 599–601.
273. Allan, A.M. & R.A. Harris. 1986. Gamma-aminobutyric acid agonists and antagonists alter chloride flux across brain membranes. *Mol. Pharmacol.* **29**: 497–505.
274. Mihic, S.J., Q. Ye, M.J. Wick, *et al.* 1997. Sites of alcohol and volatile anaesthetic action on GABA<sub>A</sub> and glycine receptors. *Nature* **389**: 385–389.
275. Rossow, I. 2001. Alcohol and homicide: a cross-cultural comparison of the relationship in 14 European countries. *Addiction* **96**: S77–S92.
276. Rossow, I. 2004. Alcohol consumption and homicides in Canada, 1950–1999. *Contemp. Drug Probl.* **31**: 541–559.
277. Bye, E.K. 2008. Alcohol and homicide in Eastern Europe: A time series analysis of six countries. *Homicide Stud.* **12**: 7–27.
278. Landberg, J. & T. Norström. 2011. Alcohol and homicide in Russia and the United States: a comparative analysis. *J. Stud. Alcohol Drugs* **72**: 723–730.
279. Norström, T. 2011. Alcohol and homicide in the United States: is the link dependent on wetness? *Drug Alcohol Rev.* **30**: 458–465.